

Product datasheet

Anti-ND4 antibody [9E4-2D8] - N-terminal ab219822

1 Image

Overview

Product name	Anti-ND4 antibody [9E4-2D8] - N-terminal
Description	Mouse monoclonal [9E4-2D8] to ND4 - N-terminal
Host species	Mouse
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide corresponding to Human ND4 (N terminal). Database link: P03905
Positive control	WB: Mitochondria from cultured normal control human dermal fibroblasts neonatal (HDFn).
General notes	<u>Western blot advice:</u> Hydrophobic intrinsic membrane proteins such as the core mtDNA-encoded proteins of the mitochondrial OXPHOS complexes tend to run faster in SDS-PAGE than predicted by their amino acid composition. This is likely due to incomplete unfolding of the protein and a more negative charge:mass ratio.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 0.36% HEPES, 0.87% Sodium chloride
Purity	Protein L purified
Purification notes	Purified from hybridoma cell culture supernatant by Protein L affinity chromatography from fetal bovine serum containing medium (Protein L does not bind bovine IgG).
Clonality	Monoclonal
Clone number	9E4-2D8
Isotype	IgG2a
Light chain type	kappa

Applications

Our [Abpromise guarantee](#) covers the use of **ab219822** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use a concentration of 4 µg/ml. Detects a band of approximately 37 kDa (predicted molecular weight: 52 kDa). Western blot using whole cell extracts is not recommended.

Target

Function Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

Involvement in disease Defects in MT-ND4 are a cause of Leber hereditary optic neuropathy (LHON) [MIM:535000]. LHON is a maternally inherited disease resulting in acute or subacute loss of central vision, due to optic nerve dysfunction. Cardiac conduction defects and neurological defects have also been described in some patients. LHON results from primary mitochondrial DNA mutations affecting the respiratory chain complexes.

Defects in MT-ND4 are a cause of Leber hereditary optic neuropathy with dystonia (LDYT) [MIM:500001]; also called familial dystonia with visual failure and striatal lucencies. LDYT is part of a spectrum of Leber hereditary optic neuropathy. It is characterized by the association of optic atrophy and central vision loss with dystonia.

Defects in MT-ND4 are a cause of mitochondrial encephalomyopathy with lactic acidosis and stroke-like episodes syndrome (MELAS) [MIM:540000]. MELAS is a genetically heterogenous disorder, characterized by episodic vomiting, seizures, and recurrent cerebral insults resembling strokes and causing hemiparesis, hemianopsia, or cortical blindness.

Sequence similarities Belongs to the complex I subunit 4 family.

Cellular localization Mitochondrion membrane.

Images



Western blot - Anti-ND4 antibody [9E4-2D8] - N-terminal (ab219822)

All lanes : Anti-ND4 antibody [9E4-2D8] - N-terminal (ab219822) at 4 µg/ml

Lane 1 : Mitochondria from cultured normal control human dermal fibroblasts neonatal (HDFn)

Lane 2 : Mitochondria from HDFn cells depleted of mtDNA by long-term proliferation in the presence of ethidium bromide

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP-labeled Goat-anti-mouse IgG

Developed using the ECL technique.

Predicted band size: 52 kDa

Observed band size: 37 kDa

[why is the actual band size different from the predicted?](#)

Mitochondrial proteins solubilized in 2% SDS were separated by SDS-PAGE and then transferred to PVDF membranes in CAPS buffer.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors