

Product datasheet

Anti-Nanog antibody [EPR2027(2)] ab109250

Recombinant RabMAb

★★★★☆ 11 Abreviews 111 References 14 Images

Overview

Product name	Anti-Nanog antibody [EPR2027(2)]
Description	Rabbit monoclonal [EPR2027(2)] to Nanog
Host species	Rabbit
Specificity	100% identities with NANOGP8
Tested applications	Suitable for: WB, IHC-P, ICC/IF, Flow Cyt (Intra), IP
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: NCCIT cell lysate. IHC-P: Human seminoma tissue, Human dysgerminoma tissue and Human embryonal carcinoma tissue. ICC/IF: Human embryonic carcinoma and Human liver cell lines. IP: NCCIT whole cell lysate.
General notes	<p>Please note that Nanog is expressed variably in different tissues and that optimisation may be required depending on the tissue used for the experiment.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p> <p>Mouse: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 59% PBS, 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR2027(2)
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab109250 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★ (4)	1/1000 - 1/10000. Detects a band of approximately 37 kDa (predicted molecular weight: 35 kDa).
IHC-P	★★★★☆ (3)	1/100 - 1/250. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols . Antigen retrieval is recommended.
ICC/IF	★★★★☆ (4)	1/100 - 1/250.
Flow Cyt (Intra)		1/70. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IP		1/40.

Target

Function	Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophoctoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes (By similarity). Acts as a transcriptional activator or repressor (By similarity). Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3' (By similarity). When overexpressed, promotes cells to enter into S phase and proliferation.
Tissue specificity	Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.

Sequence similarities

Belongs to the Nanog homeobox family.
Contains 1 homeobox DNA-binding domain.


Developmental stage

Expressed in embryonic stem (ES) and carcinoma (EC) cells. Expressed in inner cell mass (ICM) of the blastocyst and gonocytes between 14 and 19 weeks of gestation (at protein level). Not expressed in oocytes, unfertilized oocytes, 2-16 cell embryos and early morula (at protein level). Expressed in embryonic stem cells (ES). Expression decreases with ES differentiation.

Cellular localization

Nucleus.

Images



All lanes : Anti-Nanog antibody [EPR2027(2)] (ab109250) at 1/2000 dilution

Lane 1 : NCCIT (Human pluripotent embryonic carcinoma epithelial cell) whole cell lysates with 5% NFDN/TBST

Lane 2 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates with 5% NFDN/TBST

Lane 3 : HEK-293 (Human embryonic kidney epithelial cell) whole cell lysates with 5% NFDN/TBST

Lane 4 : MDA-MB-231 (Human breast adenocarcinoma epithelial cell) whole cell lysates with 5% NFDN/TBST

Lane 5 : HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates with 5% NFDN/TBST

Lane 6 : Huh7 (Human hepatocellular carcinoma epithelial cell) whole cell lysates with 5% NFDN/TBST

Lane 7 : HCT 116 (Human colorectal carcinoma epithelial cell) whole cell lysates with 5% NFDN/TBST

Lane 8 : PANC-1 (Human pancreatic epithelioid carcinoma epithelial cell) whole cell lysates with 5% NFDN/TBST

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 35 kDa

Observed band size: 37 kDa


Exposure time: 30 seconds

Nanog is highly expressed in cancer stem cells. Although some papers support the expression in undifferentiated cancer cell lines, such as HeLa (PMID: 22337995, 28092370), MDA-MB-231 (PMID: 28401007, 25919570), HepG2 (PMID: 29477378), Huh7 (PMID: 26919045), HCT 116 (PMID: 25249558, 28092370) and PANC-1 (PMID: 28703793, 25846752), ab109250 can't detect the target band in these cell lines, even at the dilution of 1:200.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human seminoma tissue labelling Nanog with purified ab109250 at 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a goat anti-rabbit IgG H&L (HRP) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Nanog antibody [EPR2027(2)] (ab109250)



ab109250 (purified) at 1/40 dilution (1.5 µg/ml) immunoprecipitating Nanog in NCCIT whole cell lysate.

Lane 1 (input): NCCIT(Human pluripotent embryonic carcinoma epithelial cell) whole cell lysate 10µg


Lane 2 (+): ab109250 & NCCIT whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab109250 in NCCIT whole cell lysate

For western blotting, ab109250 at 1/500 dilution (1.5 µg/ml) VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used for detection at 1/1000 dilution.

Blocking and diluting buffer: 5% NFD /TBST.

Immunoprecipitation - Anti-Nanog antibody [EPR2027(2)] (ab109250)




Western blot - Anti-Nanog antibody [EPR2027(2)]
(ab109250)

Anti-Nanog antibody [EPR2027(2)] (ab109250) at 1/1000 dilution
(unpurified) + NCCIT cell lysate at 10 µg


Predicted band size: 35 kDa

Observed band size: 37 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-Nanog antibody
[EPR2027(2)] (ab109250)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded
sections) analysis of Human seminoma tissue labelling Nanog with
unpurified ab109250 at 1/100.




Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-Nanog antibody
[EPR2027(2)] (ab109250)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded
sections) analysis of Human dysgerminoma tissue labelling Nanog
with unpurified ab109250 at 1/100.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human embryonal carcinoma tissue labelling Nanog with unpurified ab109250 at 1/100.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Nanog antibody [EPR2027(2)] (ab109250)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human normal colon tissue shows negative staining of Nanog with unpurified ab109250.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Nanog antibody [EPR2027(2)] (ab109250)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Nanog antibody [EPR2027(2)] (ab109250)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human adult kidney tissue shows negative staining of Nanog with unpurified ab109250.



Flow Cytometry (Intracellular) - Anti-Nanog antibody [EPR2027(2)] (ab109250)


Intracellular Flow Cytometry analysis of NCCIT cells labelling Nanog with purified ab109250 at 1/70 (red). Cells were fixed with 4% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.



Immunocytochemistry/ Immunofluorescence - Anti-Nanog antibody [EPR2027(2)] (ab109250)


This image is courtesy of an anonymous Abreview

Immunocytochemistry/Immunofluorescence analysis of Human Liver cells labelling Nanog with unpurified ab109250. Cells were fixed with Paraformaldehyde, permeabilized with Triton X-100 0.1% and blocked with 1% BSA for 12 hours at 4°C. Sample was incubated with primary antibody (1/500 in PBS) for 16 hour at 4°C. An Alexa Fluor[®]647-conjugated Donkey anti-rabbit(1/1000) IgG polyclonal was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-Nanog antibody [EPR2027(2)] (ab109250)

Immunocytochemistry/Immunofluorescence analysis of embryonic carcinoma cells labelling Nanog with unpurified ab109250 at 1/100.



Immunocytochemistry/ Immunofluorescence - Anti-Nanog antibody [EPR2027(2)] (ab109250)

Immunocytochemistry/Immunofluorescence analysis of NCCIT(human pluripotent embryonal carcinoma) cells labelling Nanog with purified ab109250 at 1/250. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. [ab150077](#), an Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. [ab7291](#), a mouse anti-tubulin (1/1000) and [ab150120](#), an Alexa Fluor[®] 594-conjugated goat anti-mouse IgG (1/1000) were also used.

Control 1: primary antibody (1/100) and secondary antibody, [ab150120](#), an Alexa Fluor[®] 594-conjugated goat anti-mouse IgG (1/500).

Control 2: [ab7291](#) (1/1000) and secondary antibody, [ab150077](#), an Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/500).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Nanog antibody [EPR2027(2)] (ab109250)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors