

Product datasheet

Anti-Neuroglycan C antibody ab31946

★★★★☆ 1 Abreviews 1 References 4 Images

Overview

Product name	Anti-Neuroglycan C antibody
Description	Rabbit polyclonal to Neuroglycan C
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, IHC-FoFr, WB
Species reactivity	Reacts with: Mouse, Rat Predicted to work with: Cow, Human
Immunogen	Synthetic peptide conjugated to KLH derived from within residues 450 - 550 of Mouse Neuroglycan C. Read Abcam's proprietary immunogen policy (Peptide available as ab32755 .)
Positive control	This antibody gave a positive signal in the following lysates: Spinal Cord (Mouse) Tissue Lysate Spinal Cord (Rat) Lysate. Brain (Mouse) Tissue Lysate Brain (Rat) Tissue Lysate

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab31946** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		

Application	Abreviews	Notes
IHC-FoFr	★★★★☆	

WB

Application notes

ICC/IF: Use at a concentration of 1 µg/ml.
 IHC-FoFr: 1/300 (See abreview for more details)
 WB: Use at a concentration of 1 µg/ml. Detects a band of approximately 60,45 kDa (predicted molecular weight: 60,58,48 kDa).

Not yet tested in other applications.
 Optimal dilutions/concentrations should be determined by the end user.

Target

Function

May function as a growth and differentiation factor involved in neuritogenesis. May induce ERBB3 activation.

Tissue specificity

Restricted to brain (at protein level).

Sequence similarities

Contains 1 EGF-like domain.

Developmental stage

Expressed in brain of 3 months, 5 and 10-year-old individuals.


Post-translational modifications

N-glycosylated.
 O-glycosylated; contains chondroitin sulfate glycans. Part-time proteoglycan, expressed in part as a proteoglycan exhibiting chondroitin sulfate glycans and in part as a non-proteoglycan form. The relative amount of both forms depends on tissues and tissues maturation.
 Phosphorylated; in intracellular and extracellular parts.

Cellular localization

Cell membrane. Endoplasmic reticulum membrane. Golgi apparatus membrane. In neurons, localizes to synaptic junctions (By similarity). Also detected in the endoplasmic reticulum and the Golgi (By similarity). Partially enriched in lipid rafts.


Images



Immunohistochemistry (PFA perfusion fixed frozen sections) - Anti-Neuroglycan C antibody (ab31946)

ab31946 staining Neuroglycan C in rat brain tissue section by Immunohistochemistry (PFA perfusion fixed frozen sections). Tissue from 4% PFA perfused animals underwent overnight fixation in 4% paraformaldehyde, cryoprotected in 30% sucrose and cut using cryostat. The primary antibody was diluted, 1/300 (PBS + 0.3% Triton X100) and incubated with sample for 18 hours at 20°C. An Alexa Fluor® 488 conjugated goat polyclonal to rabbit IgG was used at 1/1000 dilution, as secondary.

This image is courtesy of an Abreview submitted by Dr Sophie Pezet.



Western blot - Anti-Neuroglycan C antibody (ab31946)

All lanes : Anti-Neuroglycan C antibody (ab31946) at 1 µg/ml

Lane 1 : Mouse Spinal Cord Lysate

Lane 2 : Rat Spinal Cord Lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : IRDye 680 Conjugated Goat Anti-Rabbit IgG (H+L) at 1/15000 dilution


Performed under reducing conditions.

Predicted band size: 60,58,48 kDa

Observed band size: 45,60 kDa

[why is the actual band size different from the predicted?](#)

It is impossible to conclude from the data whether the higher observed band is Neuroglycan C isoform 3 (60 kDa) or Neuroglycan C isoform 1 (57 kDa). The epitope for this antibody is present in all 3 isoforms of the Neuroglycan C protein. We suspect the lower band is Isoform 2.



Western blot - Anti-Neuroglycan C antibody (ab31946)

All lanes : Anti-Neuroglycan C antibody (ab31946) at 1 µg/ml

Lane 1 : Brain (Mouse) Tissue Lysate

Lane 2 : Brain (Rat) Tissue Lysate - normal tissue

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : IRDye 680 Conjugated Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution

Performed under reducing conditions.

Predicted band size: 60,58,48 kDa

Observed band size: 45 kDa [why is the actual band size different from the predicted?](#)



Immunocytochemistry/ Immunofluorescence - Anti-Neuroglycan C antibody (ab31946)

ICC/IF image of ab31946 stained human HeLa cells. The cells were methanol fixed (5 min), permeabilised in TBS-T (20 min) and incubated with the antibody (ab31946, 1µg/ml) for 1h at room temperature. 1%BSA / 10% normal goat serum / 0.3M glycine was used to quench autofluorescence and block non-specific protein-protein interactions. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors