

Product datasheet

Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224] ab75749

KO VALIDATED Recombinant RabMAB

7 References 5 Images

Overview

Product name	Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224]
Description	Rabbit monoclonal [EPR3224] to PACT (PKR activating protein) / PRKRA
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P Unsuitable for: Flow Cyt, ICC or IP
Species reactivity	Reacts with: Human Predicted to work with: Mouse, Rat
Immunogen	Synthetic peptide within Human PACT (PKR activating protein)/ PRKRA aa 300-400 (C terminal). The exact sequence is proprietary.
Positive control	WB: HEK-293T, K562, Jurkat and HepG2 whole cell lysate (ab7900). IHC-P: Human brain tissue.
General notes	

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAB[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAB[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture

	supernatant
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR3224
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab75749 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		
IHC-P		

Application notes

IHC-P: 1/100 - 1/250. Perform heat mediated antigen retrieval using 0.01M Sodium Citrate Buffer, pH 6.0 before commencing with IHC staining protocol.
 WB: 1/1000 - 1/5000. Detects a band of approximately 34 kDa (predicted molecular weight: 34 kDa).

Is unsuitable for Flow Cyt, ICC or IP.

Not yet tested in other applications.
 Optimal dilutions/concentrations should be determined by the end user.

Target

Function

Activates EIF2AK2/PKR in the absence of double stranded RNA (dsRNA), leading to phosphorylation of EIF2S1/EIF2-alpha and inhibition of translation and induction of apoptosis. Required for siRNA production by DICER1 and for subsequent siRNA-mediated post-transcriptional gene silencing. Does not seem to be required for processing of pre-miRNA to miRNA by DICER1.

Involvement in disease

Defects in PRKRA are the cause of dystonia type 16 (DYT16) [MIM:612067]. DYT16 is an early-onset dystonia-parkinsonism disorder. Dystonia is defined by the presence of sustained involuntary muscle contraction, often leading to abnormal postures. DYT16 patients have progressive, generalized dystonia with axial muscle involvement, oro-mandibular (sardonic smile) and laryngeal dystonia and, in some cases, parkinsonian features.

Sequence similarities

Belongs to the PRKRA family.
 Contains 3 DRBM (double-stranded RNA-binding) domains.

Domain

Self-association may occur via interactions between DRBM domains as follows: DRBM 1/DRBM 1, DRBM 1/DRBM 2, DRBM 2/DRBM 2 or DRBM 3/DRBM3.

Post-translational modifications


Phosphorylated at Ser-246 in unstressed cells and at Ser-287 in stressed cells. Phosphorylation at Ser-246 appears to be a prerequisite for subsequent phosphorylation at Ser-287. Phosphorylation at Ser-246 and Ser-287 are necessary for activation of EIF2AK2/PKR under

conditions of stress.

Cellular localization

Cytoplasm > perinuclear region.

Images



All lanes : Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224] (ab75749) at 1/1000 dilution

Lane 1 : Wild-type HEK-293T cell lysate

Lane 2 : PRKRA knockout HEK-293T cell lysate

Lane 3 : K-562 cell lysate

Lane 4 : HepG2 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) at 1/10000 dilution

Predicted band size: 34 kDa

Observed band size: 36 kDa

Lanes 1-4: Merged signal (red and green). Green - ab75749 observed at 36 kDa. Red - loading control ab8245 observed at 36 kDa.


ab75749 Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224] was shown to specifically react with PACT in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line ab266806 (knockout cell lysate ab258141) was used. Wild-type and PACT knockout samples were subjected to SDS-PAGE. ab75749 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) were incubated at room temperature for 2.5 hours at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224] (ab75749)

ab75749 at 1/100 dilution staining PACT (PKR activating protein) / PRKRA in paraffin-embedded human brain tissue.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224] (ab75749)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: PACT (PKR activating protein)/PRKRA knockout HAP1 cell lysate (20 µg)

Lane 3: K562 cell lysate (20 µg)

Lane 4: HepG2 cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab75749 observed at 36 kDa. Red - loading control, [ab18058](#), observed at 124 kDa.

ab75749 was shown to specifically react with PACT (PKR activating protein)/PRKRA when PACT (PKR activating protein)/PRKRA knockout samples were used. Wild-type and PACT (PKR activating protein)/PRKRA knockout samples were subjected to SDS-PAGE. ab75749 and [ab18058](#) (loading control to Vinculin) were diluted at 1/1000 and 1/10 000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) [ab216776](#) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



All lanes : Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224] (ab75749) at 1/1000 dilution

Lane 1 : K562 cell lysate

Lane 2 : Jurkat cell lysate

Lane 3 : HepG2 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 34 kDa

Observed band size: 34 kDa

Why choose a recombinant antibody?

<p>Research with confidence Consistent and reproducible results</p>	<p>Long-term and scalable supply Recombinant technology</p>
<p>Success from the first experiment Confirmed specificity</p>	<p>Ethical standards compliant Animal-free production</p>

Anti-PACT (PKR activating protein) / PRKRA antibody [EPR3224] (ab75749)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors