


## Product datasheet

# Anti-PARN antibody ab125185

[1 References](#) [4 Images](#)

### Overview

---

<b>Product name</b>	Anti-PARN antibody
<b>Description</b>	Rabbit polyclonal to PARN
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> WB, IP, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Human <b>Predicted to work with:</b> Chimpanzee, Rhesus monkey, Orangutan 
<b>Immunogen</b>	Synthetic peptide, corresponding to a region within amino acids 589-639 of Human PARN (NP_002573.1).
<b>Positive control</b>	WB: HeLa, Jurkat and 293T whole cell lysates. IP: HeLa cell lysate. IHC-P: Human breast carcinoma and mouse renal carcinoma tissues.
<b>General notes</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
<b>Storage buffer</b>	pH: 7 Preservative: 0.09% Sodium azide Constituent: 99% Tris citrate/phosphate
<b>Purity</b>	pH 7 to 8 Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

## Applications

**The Abpromise guarantee** Our [Abpromise guarantee](#) covers the use of ab125185 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/2000 - 1/10000. Predicted molecular weight: 73 kDa.
IP		Use at 2-10 µg/mg of lysate.
IHC-P		1/500 - 1/2000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

## Target

### Function

3'-exoribonuclease that has a preference for poly(A) tails of mRNAs, thereby efficiently degrading poly(A) tails. Exonucleolytic degradation of the poly(A) tail is often the first step in the decay of eukaryotic mRNAs and is also used to silence certain maternal mRNAs translationally during oocyte maturation and early embryonic development. Interacts with both the 3'-end poly(A) tail and the 5'-end cap structure during degradation, the interaction with the cap structure being required for an efficient degradation of poly(A) tails. Involved in nonsense-mediated mRNA decay, a critical process of selective degradation of mRNAs that contain premature stop codons. Also involved in degradation of inherently unstable mRNAs that contain AU-rich elements (AREs) in their 3'-UTR, possibly via its interaction with KHSRP. Probably mediates the removal of poly(A) tails of AREs mRNAs, which constitutes the first step of destabilization.

### Tissue specificity

Ubiquitous.


### Sequence similarities

Belongs to the CAF1 family.  
Contains 1 R3H domain.

### Cellular localization

Nucleus. Cytoplasm. Nucleus > nucleolus. Some nuclear fraction is nucleolar.

## Images



**All lanes** : Anti-PARN antibody (ab125185) at 0.1 µg/ml

**Lane 1** : HeLa whole cell lysate at 50 µg

**Lane 2** : HeLa whole cell lysate at 15 µg


**Lane 3** : 293T whole cell lysate at 50 µg

**Lane 4** : Jurkat whole cell lysate at 50 µg

Developed using the ECL technique.


**Predicted band size:** 73 kDa

**Exposure time:** 3 minutes



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PARN antibody (ab125185)

Immunohistochemical analysis of formalin-fixed, paraffin-embedded human breast carcinoma tissue labeling PARN with ab125185 at a 1/1000 dilution.




Immunoprecipitation - Anti-PARN antibody (ab125185)

ab125185, at 6  $\mu\text{g}/\text{mg}$  lysate, detecting PARN in HeLa cell lysate (1 mg of immunoprecipitate, 20% loaded in the lane) by Immunoprecipitation.

For the subsequent Western blot, ab125185 was used at 1  $\mu\text{g}/\text{ml}$ .

The blot was developed using the ECL technique, with an exposure time of 3 seconds.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PARN antibody (ab125185)

Immunohistochemical analysis of formalin-fixed, paraffin-embedded mouse renal carcinoma tissue labeling PARN with ab125185 at a 1/1000 dilution.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors