


Product datasheet

Anti-Paxillin (phospho S126) antibody ab24402

1 References 4 Images

Overview

<b>Product name</b>	Anti-Paxillin (phospho S126) antibody
<b>Description</b>	Rabbit polyclonal to Paxillin (phospho S126)
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> IHC-P, ICC/IF, WB
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Human <b>Predicted to work with:</b> Chicken 
<b>Immunogen</b>	Synthetic peptide corresponding to Human Paxillin (phospho S126).
<b>Positive control</b>	WB: RAW 264.7 cells treated with LPS. A549. HepG2 IHC-P: human breast carcinoma ICC/IF: NIH/3T3
<b>General notes</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>Storage buffer</b>	pH: 7.30 Preservative: 0.05% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.1% BSA
<b>Purity</b>	Immunogen affinity purified
<b>Purification notes</b>	Purified from rabbit serum by sequential epitope-specific chromatography. The antibody has been negatively preadsorbed using a non-phosphopeptide corresponding to the site of phosphorylation to remove antibody that is reactive with non-phosphorylated paxillin. The final product is generated by affinity chromatography using a paxillin-derived peptide that is phosphorylated at serine 126.

Clonality	Polyclonal
Isotype	IgG

## Applications

**The Abpromise guarantee** Our [Abpromise guarantee](#) covers the use of ab24402 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		1/20. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		1/250.
WB		1/1000. Detects a band of approximately 68 kDa.

## Target


**Function** Cytoskeletal protein involved in actin-membrane attachment at sites of cell adhesion to the extracellular matrix (focal adhesion).

**Sequence similarities** Belongs to the paxillin family.  
Contains 4 LIM zinc-binding domains.

**Post-translational modifications** Phosphorylated on tyrosine residues during integrin-mediated cell adhesion, embryonic development, fibroblast transformation and following stimulation of cells by mitogens.

**Cellular localization** Cytoplasm > cytoskeleton. Cell junction > focal adhesion.

## Images



Western blot - Anti-Paxillin (phospho S126) antibody (ab24402)

**All lanes** : Anti-Paxillin (phospho S126) antibody (ab24402) at 1/1000 dilution

**Lane 1** : Lysate from untreated RAW 264.7 cells

**Lane 2** : Lysate from RAW 264.7 cells treated with 1 µg/mL LPS for 60 minutes

**Lane 3** : Lysate from RAW 264.7 cells treated with 1 µg/mL LPS for 60 minutes with non-phosphorylated peptide corresponding to the immunogen

**Lane 4** : Lysate from RAW 264.7 cells treated with 1 µg/mL LPS for 60 minutes with generic phosphoserine-containing peptide

**Lane 5** : Lysate from RAW 264.7 cells treated with 1 µg/mL LPS for 60 minutes with phosphopeptide corresponding to Paxillin [pS<sup>91</sup>]

**Lane 6** : Lysate from RAW 264.7 cells treated with 1 µg/mL LPS


for 60 minutes with phosphopeptide immunogen

### Secondary

**All lanes** : Goat F(ab)<sub>2</sub> anti-rabbit IgG HRP conjugate.

**Observed band size:** 68 kDa

Lysates were resolved on a 10% polyacrylamide gel and transferred to PVDF. The membrane was blocked with a 5% milk-TBST buffer for one hour at room temperature, and then incubated with the Paxillin [pS<sup>126</sup>] antibody for two hours at room temperature in a 1% milk-TBST buffer, following its prior incubation with or without peptides. Following incubation with goat F(ab')<sub>2</sub> anti-rabbit IgG HRP conjugate and bands were detected using the Pierce SuperSignal method. The data show that only the phosphopeptide corresponding to Paxillin [pS 126 ] blocks the antibody signal, thereby demonstrating the specificity of the antibody. No competition was seen following incubation with paxillin phosphopeptides to S<sup>130</sup>, S<sup>178</sup>, S<sup>380</sup>, S<sup>455</sup>, or S<sup>479</sup> (not shown). The antibody was also shown to be specific using 293 cells transfected with wild-type EGFP-tagged human paxillin treated with EGF (not shown).



Western blot - Anti-Paxillin (phospho S126) antibody (ab24402)

**All lanes** : Anti-Paxillin (phospho S126) antibody (ab24402) at 1/1000 dilution

**Lane 1** : A549 (Human lung carcinoma cell line) whole cell lysate

**Lane 2** : A549 whole cell lysate treated with 0.1 ug/mL of HGF treatment for 10 minutes

**Lane 3** : HepG2 (Human liver hepatocellular carcinoma cell line) whole cell lysate

**Lane 4** : HepG2 whole cell lysate with 0.1 ug/mL of HGF treatment for 10 minutes


Lysates/proteins at 30 µg per lane.

### Secondary

**All lanes** : Goat Anti-Rabbit IgG (H+L) Secondary Antibody, HRP conjugate at 1/1000 dilution

**Observed band size:** 68 kDa


The membrane was probed with the relevant primary and secondary Antibody following blocking with 5 % skimmed milk. Chemiluminescent detection was performed using Pierce™ ECL Western Blotting Substrate



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Paxillin (phospho S126) antibody (ab24402)

Immunohistochemical analysis of human breast carcinoma labeling Paxillin (phospho S126) with ab24402 at 1/20 dilution (right) compared to a negative control without primary antibody (left).

To expose target proteins, antigen retrieval was performed using 10mM sodium citrate (pH 6.0), microwaved for 8-15 min. Following antigen retrieval, tissues were blocked in 3% H<sub>2</sub>O<sub>2</sub>-methanol for 15 min at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with ab24402 diluted in 3% BSA-PBS at a dilution of 1/20 overnight at 4°C in a humidified chamber. Tissues were washed extensively in PBST and detection was performed using an HRP-conjugated secondary antibody followed by colorimetric detection using a DAB kit. Tissues were counterstained with hematoxylin and dehydrated with ethanol and xylene to prep for mounting



Immunocytochemistry/ Immunofluorescence - Anti-Paxillin (phospho S126) antibody (ab24402)

Immunofluorescence analysis of 70% confluent log phase NIH/3T3 cells treated with 0.5 µg of PDGF for 10 minutes. The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 10 minutes, and blocked with 1% BSA for 1 hour at room temperature. Labeling Paxillin (phospho S126) with ab24402 at 1/250 dilution in 0.1% BSA and incubated for 3 hours at room temperature. Followed by Goat anti-Rabbit IgG (H+L) Superclonal™ Secondary Antibody, Alexa Fluor® 488 conjugate at a dilution of 1/2000 for 45 minutes at room temperature (Panel a: green). Nuclei (Panel b: blue) were stained with SlowFade® Gold Antifade Mountant with DAPI. F-actin (Panel c: red) was stained with Rhodamine Phalloidin at 1/300 dilution. Panel d is a merged image showing punctuated nuclear localization. Panel e is untreated cell with no signal. Panel f is a no primary antibody control. The images were captured at 60X magnification.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors