

Product datasheet

Anti-POMC antibody [EPR22534-165] ab254257

Recombinant RabMAb

★★★★☆ [1 Abreviews](#) [1 References](#) [11 Images](#)

Overview

Product name	Anti-POMC antibody [EPR22534-165]
Description	Rabbit monoclonal [EPR22534-165] to POMC
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P, IHC-Fr, ICC/IF Unsuitable for: IP
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: Mouse, human and rat pituitary tissue lysate. IHC-P: Mouse pituitary and spleen; human and rat pituitary tissue. IHC-Fr: Mouse and rat pituitary and liver tissue. ICC/IF: AtT-20 cells.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR22534-165

Isotype

IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab254257 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		1/1000. Detects a band of approximately 35 kDa (predicted molecular weight: 29 kDa).
IHC-P		1/4000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IHC-Fr	★★★★★ (1)	1/500.
ICC/IF		1/100.

Application notes

Is unsuitable for IP.

Target

Images



Western blot - Anti-POMC antibody [EPR22534-165] (ab254257)

Lanes 1-4 : Anti-POMC antibody [EPR22534-165] (ab254257) at 1/5000 dilution

Lanes 5-6 : Anti-POMC antibody [EPR22534-165] (ab254257) at 1/1000 dilution

Lane 1 : Mouse pituitary lysate

Lane 2 : Mouse liver lysate

Lane 3 : Human pituitary lysate

Lane 4 : Human liver lysate

Lane 5 : Rat pituitary lysate

Lane 6 : Rat liver lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 29 kDa

Observed band size: 35, 16 kDa

Exposure time:

Lanes 1-2: 3 seconds;


Lanes 3-4: 10 seconds;

Lanes 5-6: 32 seconds

Blocking buffer and concentration: 5% NFDm/TBST

Negative control: liver (lanes 2, 4 and 6, PMID: 10395956)

POMC is a glycoprotein and can be processed into 16kDa N-POMC. The molecular weight observed is consistent with what has been described in the literature (PMID 1814247, PMID 26880796, PMID 20639461).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-POMC antibody [EPR22534-165] (ab254257)

Immunohistochemical analysis of paraffin-embedded human pituitary tissue labeling POMC with ab254257 at 1/4000 dilution at room temperature for 30 mins, followed by ready to use rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)). Positive staining on human pituitary is observed (PMID: 21622576).

Counterstained with hematoxylin. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins was performed. The immunostaining was performed on a Leica Biosystems BOND[®] RX instrument.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is ready to use rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-POMC antibody [EPR22534-165] (ab254257)


Negative control: No staining on mouse spleen (PMID: 10395956)

Immunohistochemical analysis of paraffin-embedded mouse pituitary tissue labeling POMC with ab254257 at 1/4000 dilution at room temperature for 30 mins, followed by ready to use rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)). Counterstained with hematoxylin. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins was performed. The immunostaining was performed on a Leica Biosystems BOND[®] RX instrument.



Immunohistochemistry (Frozen sections) - Anti-POMC antibody [EPR22534-165] (ab254257)


Immunohistochemical analysis of 0.2% Triton X-100 permeabilized, 4% PFA fixed mouse pituitary tissue labeling POMC with ab254257 at 1/500 dilution, followed by AlexaFluor[®]488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution. DAPI was used as nuclear counterstain. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20) was performed. Positive staining on mouse pituitary (PMID: 18601691). Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is AlexaFluor[®]488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution



Immunohistochemistry (Frozen sections) - Anti-POMC antibody [EPR22534-165] (ab254257)

Negative control: No stain on liver (PMID: 14578285)


Immunohistochemical analysis of 0.2% Triton X-100 permeabilized, 4% PFA fixed mouse liver tissue labeling POMC with ab254257 at 1/100 dilution, followed by AlexaFluor[®]488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution. DAPI was used as nuclear counterstain. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20) was performed. No staining on mouse liver. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is AlexaFluor[®]488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-POMC antibody [EPR22534-165] (ab254257)

Immunohistochemical analysis of paraffin-embedded rat pituitary tissue labeling POMC with ab254257 at 1/4000 dilution at room temperature for 30 mins, followed by ready to use rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)). Positive staining on rat pituitary is observed (PMID: 21622576). Counterstained with hematoxylin. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins was performed. The immunostaining was performed on a Leica Biosystems BOND® RX instrument.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is ready to use rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-POMC antibody [EPR22534-165] (ab254257)

Immunohistochemical analysis of paraffin-embedded mouse pituitary tissue labeling POMC with ab254257 at 1/4000 dilution at room temperature for 30 mins, followed by ready to use rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)). Positive staining on mouse pituitary is observed (PMID: 21622576). Counterstained with hematoxylin. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins was performed. The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is ready to use rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)).




Immunohistochemistry (Frozen sections) - Anti-POMC antibody [EPR22534-165] (ab254257)

Negative control: No stain on liver (PMID: 14578285)

Immunohistochemical analysis of 0.2% Triton X-100 permeabilized, 4% PFA fixed rat liver tissue labeling POMC with ab254257 at 1/100 dilution, followed by AlexaFluor®488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution. DAPI was used as nuclear counterstain. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20) was performed. No staining on rat liver.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is AlexaFluor®488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution



Immunohistochemistry (Frozen sections) - Anti-POMC antibody [EPR22534-165] (ab254257)

Immunohistochemical analysis of 0.2% Triton X-100 permeabilized, 4% PFA fixed rat pituitary tissue labeling POMC with ab254257 at 1/500 dilution, followed by AlexaFluor®488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution. DAPI was used as nuclear counterstain. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20) was performed. Positive staining on rat pituitary (PMID: 18601691).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is AlexaFluor®488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution



Immunocytochemistry/ Immunofluorescence - Anti-POMC antibody [EPR22534-165] (ab254257)

Immunocytochemical analysis of 100% methanol fixed, 0.1% TritonX-100 permeabilised AtT-20 cells labeling POMC with ab254257 at 1/100 dilution. [ab150077](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) was used as a secondary antibody at 1/1000 dilution. Counterstained with [ab195889](#) Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) at 1/200 dilution. Nuclear staining: DAPI.

Confocal image showing cytoplasmic staining in AtT-20 cell line

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-POMC antibody [EPR22534-165] (ab254257)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors