

Product datasheet

Fibroblast Marker (Vimentin, alpha smooth muscle Actin, Hsp47, S100A4) Antibody Panel - Human, Mouse ab254015

Recombinant

★★★★★ [1 Abreviews](#) [2 References](#) [11 Images](#)

Overview

Product name	Fibroblast Marker (Vimentin, alpha smooth muscle Actin, Hsp47, S100A4) Antibody Panel - Human, Mouse
Species reactivity	Reacts with: Mouse, Human
Product overview	<p>Fibroblast Marker (Vimentin, alpha smooth muscle Actin, Hsp47, S100A4) Antibody Panel - Human, Mouse ab254015 contains multiple trial-sized versions of anti-human and mouse antibody clones against Vimentin, alpha smooth muscle Actin, Hsp47, S100A4, specifically selected for high performance in various applications. This panel contains 4 recombinant rabbit monoclonal antibodies against human and mouse Vimentin, alpha smooth muscle Actin, Hsp47, S100A4. They are provided as a sampler panel to allow you to easily evaluate each in your required applications.</p> <p>For guidelines on how to use each antibody within the panel, please consult the individual datasheet for each antibody.</p> <p>Panel contains:</p> <ul style="list-style-type: none">- Rabbit monoclonal [EPR3776] to Vimentin (20 µL) ab92547- Rabbit monoclonal [E184] to alpha smooth muscle Actin (20 µL) ab32575- Rabbit monoclonal [EPR4217] to Hsp47 (20 µL) ab109117- Rabbit monoclonal [EPR14639(2)] to S100A4 (20 µL) ab197896

Notes [Explore our range of antibody sample panels](#) designed to provide you with a variety of trial-size antibodies in a convenient and cost-effective format.

Directly conjugated versions of our antibodies are available and ready to use for multicolor flow cytometry or immunocytochemistry analysis. [Carrier-free formulations](#) are also available for easy conjugation to labels of your choice. Please refer to the 'Associated products' section below.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).


Properties

Storage instructions Store at -20°C. Please refer to protocols.

Components	1 kit
ab32575 - Anti-alpha smooth muscle Actin antibody [E184]	2 x 10µl
ab109117 - Anti-Hsp47 antibody [EPR4217]	2 x 10µl
ab197896 - Anti-S100A4 antibody [EPR14639(2)]	2 x 10µl
ab92547 - Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker	2 x 10µl


Cellular localization Vimentin: Cytoplasm. alpha smooth muscle Actin: Cytoplasm > cytoskeleton. Hsp47: Endoplasmic reticulum lumen.

Images




Immunohistochemical staining of paraffin embedded human cervical carcinoma with purified [ab92547](#) at a working dilution of 1/250. The secondary antibody used is HRP goat anti-rabbit IgG H&L ([ab97051](#)) at 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Vimentin antibody [EPR3776]


Immunohistochemical staining of paraffin embedded mouse kidney with purified **ab92547** at a working dilution of 1/250. The secondary antibody used is **Goat Anti-Rabbit IgG H&L (HRP) (ab97051) secondary antibody** at 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Immunocytochemistry/ Immunofluorescence - Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker

ab92547 staining Vimentin in wild-type HAP1 cells (top panel) and VIM knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with **ab92547** at 0.5µg/ml and **ab195889** at 1/250 dilution (shown in pseudocolour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with **Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed (ab150081) secondary antibody** at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha smooth muscle Actin antibody [E184]

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human smooth muscle tissue labeling alpha smooth muscle Actin with purified **ab32575** at a dilution of 1/200. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. **ab97051**, an HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500)

Negative control using PBS instead of primary antibody.


Counterstained with hematoxylin.



Immunocytochemistry/ Immunofluorescence - Anti-alpha smooth muscle Actin antibody [E184]

Immunocytochemistry/Immunofluorescence analysis of A431 (human epidermoid carcinoma) cells labeling alpha smooth muscle Actin (green) with purified **ab32575** at 1/500. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. Cells were counterstained with **ab7291**, anti-Tubulin (mouse mAb) at 1/1000 followed by **ab150120** Alexa Fluor[®]594 goat anti-mouse secondary (1/1000). Nuclei were counterstained with DAPI (blue).

For negative control 1, rabbit primary antibody and anti-mouse secondary antibody (**ab150120**) were used. For negative control 2, **ab7291** (mouse primary antibody) was used followed by anti-rabbit secondary antibody (**ab150077**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha smooth muscle Actin antibody [E184]

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse smooth muscle tissue labeling alpha smooth muscle Actin with purified **ab32575** at a dilution of 1/200. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. **ab97051**, an HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500).


Negative control using PBS instead of primary antibody.

Counterstained with hematoxylin.



Immunocytochemistry/ Immunofluorescence - Anti-alpha smooth muscle Actin antibody [E184]

Immunocytochemistry/ Immunofluorescence analysis of NIH/3T3(Mouse embryonic fibroblast) cells labeling alpha smooth muscle Actin with purified **ab32575** at 1/500 dilution (5.2 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 dilution (2 µg/ml) dilution. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A4 antibody [EPR14639(2)]

Immunohistochemical analysis of paraffin-embedded Human gastric carcinoma tissue labeling S100A4 using **ab197896** at 1/2000 dilution. Goat Anti-Rabbit IgG H&L (HRP) **ab97051** was used as a secondary antibody at 1/500 dilution. Cells were counterstained with Hematoxylin.

Inset image: negative control obtained using PBS instead of **ab197896** and secondary antibody only.

Note: Cytoplasm and nuclear staining on human gastric carcinoma tissue was observed.



Western blot - Anti-Hsp47 antibody [EPR4217]

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: Hsp47 knockout HAP1 whole cell lysate (20 µg)

Lane 3: HeLa whole cell lysate (20 µg)

Lane 4: HepG2 whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - Anti-Hsp47 antibody [EPR4217] (**ab109117**) observed at 46 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab109117 was shown to recognize Hsp47 in wild-type HAP1 cells as signal was lost at the expected MW in Hsp47 knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and Hsp47 knockout samples were subjected to SDS-PAGE. **ab109117** and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



ab109117 staining Hsp47 in Human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed and paraffin-embedded, antigen retrieval was by heat mediation in Tris/EDTA buffer pH9. Samples were incubated with primary antibody (1/300). An undiluted HRP-conjugated mouse anti-rabbit IgG was used as the secondary antibody. Tissue counterstained with Hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp47 antibody [EPR4217]

Why choose a recombinant antibody?



Fibroblast Marker (Vimentin, alpha smooth muscle Actin, Hsp47, S100A4) Antibody Panel - Human, Mouse (ab254015)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors