



Product datasheet

5,7-Dichlorokynurenic acid, NMDA receptor glycine site antagonist ab120023

[2 Images](#)

Overview

Product name	5,7-Dichlorokynurenic acid, NMDA receptor glycine site antagonist
Description	NMDA receptor glycine site antagonist
Biological description	Potent NMDA receptor glycine site antagonist. Water soluble form available - see (ab120254).
CAS Number	131123-76-7
Chemical structure	



Properties

Chemical name	5,7-Dichloro-4-hydroxyquinoline-2-carboxylic acid
Molecular weight	258.06
Molecular formula	C ₁₀ H ₅ Cl ₂ NO ₃
PubChem identifier	1779
Storage instructions	Store at +4°C. Store under desiccating conditions. The product can be stored for up to 12 months.
Solubility overview	Soluble in 1 eq. NaOH to 50 mM
Handling	<p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Refer to SDS for further information</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked questions (FAQ) page for more details.</p>
SMILES	<chem>O=C(O)c1cc(O)c2c(Cl)cc(Cl)cc2n1</chem>
Source	Synthetic

Applications


The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab120023 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Functional Studies		Use at an assay dependent concentration.

Images



2D chemical structure image of ab120023, 5,7-Dichlorokynurenic acid, NMDA receptor glycine site antagonist



ab12416 staining cGMP in SKNSH cells treated with 5,7-Dichlorokynurenic acid (ab120023), by ICC/IF. Decrease in cGMP expression correlates with increased concentration of 5,7-Dichlorokynurenic acid, as described in literature.

The cells were incubated at 37°C for 20 minutes in media containing different concentrations of ab120023 (5,7-Dichlorokynurenic acid) in DMSO. Some samples were then further incubated with 15 μM NMDA (**ab120052**) for 5 minutes and all samples were fixed with 100% methanol for 5 minutes at -20°C and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab12416** (5 μg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team