abcam

Product datasheet

Fenofibrate, PPAR-alpha agonist ab120832

3 References 2 Images

Overview

Product nameFenofibrate, PPAR-alpha agonistDescriptionPotent, selective PPAR-α agonist

Biological description Potent and selective PPAR-α agonist (EC₅₀ values are 18 and 30 μM at mouse and human

receptors, respectively). Affinity is 10-fold less at PPAR- γ (EC $_{50}$ values are 250 and 300 μ M at

mouse and human receptors, respectively). Hypolipidemic agent.

Purity > 99%

CAS Number 49562-28-9

Chemical structure

CI CH

Properties

Chemical name 2-[4-(4-Chlorobenzoyl)phenoxy]-2-methylpropanoic acid isopropyl ester

Molecular weight 360.84

Molecular formula C₂₀H₂₁ClO₄

Storage instructions Store at Room Temperature. The product can be stored for up to 12 months.

Solubility overview Soluble in DMSO to 100 mM and in ethanol to 100 mM

Handling Wherever possible, you should prepare and use solutions on the same day. However, if you need

to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and

prior to opening the vial we recommend that you allow your product to equilibrate to room

temperature for at least 1 hour.

Refer to SDS for further information

Need more advice on solubility, usage and handling? Please visit our frequently asked

questions (FAQ) page for more details.

Source Synthetic

1


Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab120832 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Functional Studies		Use at an assay dependent concentration.

Images



2D chemical structure image of ab120832, Fenofibrate, PPARalpha agonist



HL-60 cells were incubated at 37° C for 24h with vehicle control (0 μ M) and different concentrations of fenofibrate (ab120832). Increased expression of ACADM in HL-60 cells correlates with an increase in fenofibrate concentration, as described in literature.

Whole cell lysates were prepared with RIPA buffer (containing protease inhibitors and sodium orthovanadate), 10 μ g of each were loaded on the gel and the WB was run under reducing conditions. After transfer the membrane was blocked for an hour using 5% BSA before being incubated with **ab110296** at 1 μ g/ml and **ab9484** at 1 μ g/ml overnight at 4°C. Antibody binding was detected using an anti-mouse antibody conjugated to HRP (**ab97040**) at 1/10000 dilution and visualised using ECL development solution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team