# abcam

### Product datasheet

## Sodium salicylate, anti-inflammatory agent ab120746

1 References 2 Images

Overview

Product name Sodium salicylate, anti-inflammatory agent

**Description** Cell-permeable anti-inflammatory agent

**Biological description**Cell-permeable anti-inflammatory agent. Inhibits cyclooxygenases preventing prostaglandin

formation. Inhibits components of the MAPK cascade, NF-kB and AP-1. Induces apoptosis

through p38 MAPK activation.

**Purity** > 99%

CAS Number 54-21-7

CO<sub>2</sub>Na

**Properties** 

**Chemical structure** 

Chemical name 2-Hydroxybenzoic acid sodium salt

Molecular weight160.10Molecular formula $C_7H_5NaO_3$ 

**Storage instructions** Store at Room Temperature. Store under desiccating conditions. The product can be stored for

up to 12 months.

**Solubility overview** Soluble in water to 100 mM

Handling Wherever possible, you should prepare and use solutions on the same day. However, if you need

to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and

prior to opening the vial we recommend that you allow your product to equilibrate to room

temperature for at least 1 hour.

Refer to SDS for further information

Need more advice on solubility, usage and handling? Please visit our **frequently asked** 

questions (FAQ) page for more details.

**Source** Synthetic

**Applications** 

1


#### The Abpromise quarantee

Our <u>Abpromise guarantee</u> covers the use of ab120746 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Functional Studies		Use at an assay dependent concentration.

#### **Images**



2D chemical structure image of ab120746, Sodium salicylate, antiinflammatory agent



THP1 cells were incubated at  $37^{\circ}$ C for 40h with vehicle control (0  $\mu$ M) and different concentrations of sodium salicylate (ab120746). Decreased expression of scavenging receptor SR-BI (ab52629) in THP1 cells correlates with an increase in sodium salicylate concentration, as described in literature.

Whole cell lysates were prepared with RIPA buffer (containing protease inhibitors and sodium orthovanadate), 10 µg of each were loaded on the gel and the WB was run under reducing conditions. After transfer the membrane was blocked for an hour using 5% BSA before being incubated with <a href="mailto:ab52629">ab52629</a> at 1/2000 dilution and <a href="mailto:ab52629">ab52629</a> at 1/2000 dilution and visualised using an anti-rabbit antibody conjugated to HRP (<a href="mailto:ab97051">ab97051</a>) at 1/10000 dilution and visualised using ECL development solution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team