abcam

Product datasheet

Human LDHB (Lactate Dehydrogenase B) knockout HEK-293T cell line ab255403

2 Images

Overview

Product name	Human LDHB (Lactate Dehydrogenase B) knockout HEK-293T cell line			
Parental Cell Line	HEK293T			
Organism	Human			
Mutation description	Knockout achieved by using CRISPR/Cas9, Homozygous: 1 bp insertion in exon 2			
Passage number	<20			
Knockout validation	Sanger Sequencing, Western Blot (WB)			
Tested applications	Suitable for: WB			
Biosafety level	2			
General notes	Recommended control: Human wild-type HEK293T cell line (<u>ab255449</u>). Please note a wild- type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.			
	Cryopreservation cell medium: Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.			
	Culture medium: DMEM (High Glucose) + 10% FBS			
	Initial handling guidelines: Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.			
	 Thaw the vial in 37°C water bath for approximately 1-2 minutes. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2x10⁴ cells/cm². Seeding density is given as a guide only and should be scaled to align with individual lab schedules. Incubate the culture at 37°C incubator with 5% CO₂. Cultures should be monitored daily. 			
	Subculture guidelines:			
	All seeding densities should be based on cell counts gained by established methods. A guide seeding density of 2x10 ⁴ cells/cm ² is recommended.			

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.

Cells should be passaged when they have achieved 80-90% confluence. This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our **limited use license** and **patent pages**.

We will provide viable cells that proliferate on revival.

Properties


Number of cells	1 x 10 ⁶ cells/vial, 1 mL			
Adherent /Suspension	Adherent			
Tissue	Kidney			
Cell type	epithelial			
STR Analysis	Amelogenin X D5S818: 8, 9 D13S317: 12, 14 D7S820: 11 D16S539: 9, 13 vWA: 16, 19 TH01: 7, 9.3 TPOX: 11 CSF1PO: 11, 12			
Mycoplasma free	Yes			
Storage instructions	Shipped on Dry Ice. Store in liquid nitrogen.			
Storage buffer	Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether			
Target				
Pathway	Fermentation; pyruvate fermentation to lactate; (S)-lactate from pyruvate: step 1/1.			
Involvement in disease	Note=Defects in LDHB result in deficiency of lactate dehydrogenase, a condition with no clear symptomatic consequences. Lactate dehydrogenase deficiency can probably be considered a non-disease.			
Sequence similarities	Belongs to the LDH/MDH superfamily. LDH family.			
Cellular localization	Cytoplasm.			

Applications

The Abpromise guaranteeOur Abpromise guaranteecovers the use of ab255403 in the following tested applications.The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 37 kDa.

Images	5
--------	---



Western blot - Human LDHB (Lactate Dehydrogenase B) knockout HEK293T cell line (ab255403) All lanes : Anti-Lactate Dehydrogenase B/LDH-B antibody [EP1565Y] (<u>ab53292</u>) at 1/5000 dilution

Lane 1 : Wild-type HEK-293T cell lysate Lane 2 : LDHB knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 37 kDa Observed band size: 37 kDa

Lanes 1-2: Merged signal (red and green). Green - <u>ab53292</u> observed at 37 kDa. Red - Anti-Vinculin antibody [VIN-54] observed at 124 kDa.

ab53292 was shown to react with LDHB in wild-type HEK-293T cells in western blot. Loss of signal was observed when knockout cell line ab255403 (knockout cell lysate **ab263761**) was used. Wild-type HEK-293T and LDHB knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. **ab53292** and Anti-Vinculin antibody [VIN-54] overnight at 4°C at a 1 in 5000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye[®]800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye[®]680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

Homozygous: 1 bp insertion in exon 2

Sanger Sequencing - Human LDHB knockout HEK293T cell line (ab255403)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors