

Human MAP3K2 (MEKK2) knockout A549 cell line ab267152

3 Images

Overview

Product name	Human MAP3K2 (MEKK2) knockout A549 cell line
Parental Cell Line	A549
Organism	Human
Mutation description	Knockout achieved by using CRISPR/Cas9, 1 bp insertion in exon 7 and 2 bp deletion in exon 7
Passage number	<20
Knockout validation	Sanger Sequencing, Western Blot (WB)
Tested applications	Suitable for: WB
Biosafety level	2
General notes	

Recommended control: Human wild-type A549 cell line ([ab255450](#)). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.

Cryopreservation cell medium: Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.

Culture medium: F-12K + 10% FBS

Initial handling guidelines: Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.

1. Thaw the vial in 37°C water bath for approximately 1-2 minutes.
2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.
3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2×10^3 - 1×10^4 cells/cm². Seeding density is given as a guide only and should be scaled to align with individual lab schedules.
4. Incubate the culture at 37°C incubator with 5% CO₂. Cultures should be monitored daily.

Subculture guidelines:

All seeding densities should be based on cell counts gained by established methods. A guide seeding density of 6×10^4 cells/cm² is recommended.

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.

Cells should be passaged when they have achieved 80-90% confluence.

Do not exceed 7×10^4 cells/cm².

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our [limited use license](#) and [patent pages](#).

We will provide viable cells that proliferate on revival.

Properties

Number of cells	1 x 10 ⁶ cells/vial, 1 mL
Adherent /Suspension	Adherent
Tissue	Lung
Cell type	epithelial
Disease	Carcinoma
Gender	Male
STR Analysis	Amelogenin X,YD5S818: 11 D13S317: 11 D7S820: 8, 11 D16S539: 11, 12 WWA: 14 TH01: 8,9,3 TPOX: 8,11 CSF1PO: 10, 12
Antibiotic resistance	Puromycin 1.00µg/ml
Mycoplasma free	Yes
Storage instructions	Shipped on Dry Ice. Store in liquid nitrogen.
Storage buffer	Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether

Target


Function	Component of a protein kinase signal transduction cascade. Regulates the JNK and ERK5 pathways by phosphorylating and activating MAP2K5 and MAP2K7 (By similarity). Plays a role in caveolae kiss-and-run dynamics.
Sequence similarities	Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily. Contains 1 OPR domain. Contains 1 protein kinase domain.
Post-translational modifications	Autophosphorylated.
Cellular localization	Cytoplasm. Nucleus. Upon EGF stimulation, translocates into the nucleus.

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab267152 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 70 kDa.

Images



All lanes : Anti-MEKK2 antibody [EP626Y] ([ab33918](#)) at 1/10000 dilution

Lane 1 : Wild-type A549 cell lysate

Lane 2 : MAP3K2 knockout A549 cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 70 kDa

Observed band size: 75 kDa

Lanes 1-2: Merged signal (red and green). Green - [ab33918](#) observed at 75 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) observed at 37 kDa.

[ab33918](#) was shown to react with MEKK2 in wild-type A549 cells in western blot. Loss of signal was observed when knockout cell line ab267152 (knockout cell lysate [ab257521](#)) was used. Wild-type A549 and MAP3K2 knockout A549 cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. [ab33918](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) overnight at 4°C at a 1 in 10000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

```
Mut  CTAGTAGAGATAGAAGTTCCTCCCCCAGG-ACATTCCAGATGAATTACACCAGGTTG
      |||
WT   CTAGTAGAGATAGAAGTTCCTCCCCCAGGTTACATTCCAGATGAATTACACCAGGTTG
```

Allele-1: 2 bp deletion in exon7

Sanger Sequencing - Human MAP3K2 knockout

A549 cell line (ab267152)

```
Mut  CTAGTAGAGATAGAAGTTCCTCCCCCAGGATTACATTCCAGATGAATTACACCAGGTT
      |||
WT   CTAGTAGAGATAGAAGTTCCTCCCCCAGG TTACATTCCAGATGAATTACACCAGGTT
```

Allele-2: 1 bp insertion in exon 7.

Sanger Sequencing - Human MAP3K2 knockout

A549 cell line (ab267152)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors