

Human RAB10 knockout A549 cell line ab261868

5 Images

Overview

Product name	Human RAB10 knockout A549 cell line
Parental Cell Line	A549
Organism	Human
Mutation description	Knockout achieved by CRISPR/Cas9; X = 1 bp insertion; Frameshift = 100%
Passage number	<20
Knockout validation	Immunocytochemistry (ICC), Next Generation Sequencing (NGS), Western Blot (WB)
Tested applications	Suitable for: WB, ICC
Biosafety level	1
General notes	<p>Recommended control: Human wild-type A549 cell line (ab259777). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p>Cryopreservation cell medium: Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p>Culture medium: DMEM:Hams F12 + 5% FBS</p> <p>Initial handling guidelines: Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none"> 1. Thaw the vial in 37°C water bath for approximately 1-2 minutes. 2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution. 3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2×10^3-1×10^4 cells/cm². Seeding density is given as a guide only and should be scaled to align with individual lab schedules. 4. Incubate the culture at 37°C incubator with 5% CO₂. Cultures should be monitored daily. <p>Subculture guidelines:</p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of 6×10^4 cells/cm² is recommended.</p> <p>A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.</p>

Cells should be passaged when they have achieved 80-90% confluence.
Do not exceed 7×10^4 cells/cm².

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our [limited use license](#) and [patent pages](#).

Properties

Number of cells	1 x 10 ⁶ cells/vial, 1 mL
Viability	~80%
Adherent /Suspension	Adherent
Tissue	Lung
Cell type	epithelial
Disease	Carcinoma
Gender	Male
Mycoplasma free	Yes
Storage instructions	Shipped on Dry Ice. Store in liquid nitrogen.
Storage buffer	Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether

Target

Function	<p>The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (By similarity). That Rab is mainly involved in the biosynthetic transport of proteins from the Golgi to the plasma membrane. Regulates, for instance, SLC2A4/GLUT4 glucose transporter-enriched vesicles delivery to the plasma membrane. In parallel, it regulates the transport of TLR4, a toll-like receptor to the plasma membrane and therefore may be important for innate immune response. Plays also a specific role in asymmetric protein transport to the plasma membrane within the polarized neuron and epithelial cells. In neurons, it is involved in axonogenesis through regulation of vesicular membrane trafficking toward the axonal plasma membrane while in epithelial cells, it regulates transport from the Golgi to the basolateral membrane. Moreover, may play a role in the basolateral recycling pathway and in phagosome maturation. According to PubMed:23263280, may play a role in endoplasmic reticulum dynamics and morphology controlling tubulation along microtubules and tubules fusion.</p>
Sequence similarities	Belongs to the small GTPase superfamily. Rab family.
Cellular localization	Cytoplasmic vesicle membrane. Golgi apparatus membrane. Golgi apparatus, trans-Golgi network membrane. Endosome membrane. Recycling endosome membrane. Cytoplasmic vesicle, phagosome membrane. Cell projection, cilium. Endoplasmic reticulum membrane. Associates with SLC2A4/GLUT4 storage vesicles (PubMed:22908308). Localizes to the base of the cilium (PubMed:20576682). Transiently associates with phagosomes (By similarity). Localizes to the endoplasmic reticulum at domains of new tubule growth (PubMed:23263280).

Applications


The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab261868 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration.

Images



All lanes : Anti-RAB10 antibody [MJF-R23] (**ab237703**) at 1/1000 dilution

Lane 1 : Wild-type A549 (Human lung carcinoma cell line) whole cell lysate

Lane 2 : RAB10 knockout A549 (Human lung carcinoma cell line) whole cell lysate

Lane 3 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 4 : MCF7 (Human breast adenocarcinoma cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Observed band size: 25 kDa

Lanes 1 - 4: Merged signal (red and green). Green - **ab237703** observed at 25 kDa. Red - loading control, **ab8245**, observed at 37 kDa.


ab237703 was shown to recognize RAB10 in wild-type A549 cells as signal was lost at the expected MW in RAB10 knockout cell line ab261868 (knockout cell lysate **ab261677**). Additional cross-reactive bands were observed in the wild-type and knockout samples. Wild-type and RAB10 knockout samples were subjected to SDS-PAGE. Ab237703 and **ab8245** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and

Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence -
Human RAB10 knockout A549 cell line (ab261868)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% saponin permeabilized A549 wild-type and knockout cells (ab261868) labeling RAB10 (red) with **ab237703** at 0.5 µg/ml, followed by anti-Rabbit secondary at 1/1000 dilution. The nuclear counter stain is DAPI (blue).



Western blot - Human RAB10 knockout A549 cell line (ab261868)

All lanes : Anti-RAB10 antibody [EPR13242] (**ab181367**) at 1/1000 dilution

Lane 1 : Wild-type A549 (Human lung carcinoma cell line) whole cell lysate

Lane 2 : RAB10 knockout A549 (Human lung carcinoma cell line) whole cell lysate

Lane 3 : HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 4 : MCF7 (Human breast adenocarcinoma cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Observed band size: 25 kDa

Lanes 1 -4: Merged signal (red and green). Green - **ab181367** observed at 25 kDa. Red - loading control, **ab8245**, observed at 37 kDa.


ab181367 was shown to specifically react with RAB10 in wild-type A549 cells as signal was lost in RAB10 knockout cell line

ab261868 (knockout cell lysate [ab261677](#)). Wild-type and RAB10 knockout samples were subjected to SDS-PAGE. Ab181367 and [ab8245](#) (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Human RAB10 knockout A549 cell line (ab261868)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% saponin permeabilized A549 wild-type and knockout cells (ab261868) labeling RAB10 (red) with [ab237703](#) at 0.5 µg/ml, followed by anti-Rabbit secondary at 1/1000 dilution. The nuclear counter stain is DAPI (blue).



X = 1 bp insertion

Next Generation Sequencing - Human RAB10 knockout A549 cell line (ab261868)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors