

Human SYP (Synaptophysin) knockout HEK-293T cell line ab267272

[4 Images](#)

Overview

Product name	Human SYP (Synaptophysin) knockout HEK-293T cell line
Parental Cell Line	HEK293T
Organism	Human
Mutation description	Knockout achieved by using CRISPR/Cas9, Homozygous: 8 bp deletion in exon 2
Passage number	<20
Knockout validation	Sanger Sequencing, Western Blot (WB)
Tested applications	Suitable for: WB
Biosafety level	2
General notes	<p>Recommended control: Human wild-type HEK293T cell line (ab255449). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p>Cryopreservation cell medium: Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p>Culture medium: DMEM (High Glucose) + 10% FBS</p> <p>Initial handling guidelines: Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none">1. Thaw the vial in 37°C water bath for approximately 1-2 minutes.2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2×10^4 cells/cm². Seeding density is given as a guide only and should be scaled to align with individual lab schedules.4. Incubate the culture at 37°C incubator with 5% CO₂. Cultures should be monitored daily. <p>Subculture guidelines:</p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of 2×10^4 cells/cm² is recommended.</p>

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.

Cells should be passaged when they have achieved 80-90% confluence.

This product is subject to limited use licenses from The Broad Institute and ERS Genomics Limited, and is developed with patented technology. For full details of the limited use licenses and relevant patents please refer to our [limited use license](#) and [patent pages](#).

We will provide viable cells that proliferate on revival.

Properties

Number of cells	1 x 10 ⁶ cells/vial, 1 mL
Adherent /Suspension	Adherent
Tissue	Kidney
Cell type	epithelial
STR Analysis	Amelogenin X D5S818: 8, 9 D13S317: 12, 14 D7S820: 11 D16S539: 9, 13 vWA: 16, 19 TH01: 7, 9.3 TPOX: 11 CSF1PO: 11, 12
Antibiotic resistance	Puromycin 1.00µg/ml
Mycoplasma free	Yes
Storage instructions	Shipped on Dry Ice. Store in liquid nitrogen.
Storage buffer	Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether

Target


Function	Possibly involved in structural functions as organizing other membrane components or in targeting the vesicles to the plasma membrane. Involved in the regulation of short-term and long-term synaptic plasticity.
Tissue specificity	Characteristic of a type of small (30-80 nm) neurosecretory vesicles, including presynaptic vesicles, but also vesicles of various neuroendocrine cells of both neuronal and epithelial phenotype.
Involvement in disease	Mental retardation, X-linked, SYP-related
Sequence similarities	Belongs to the synaptophysin/synaptobrevin family. Contains 1 MARVEL domain.
Domain	The calcium-binding activity is thought to be localized in the cytoplasmic tail of the protein.
Post-translational modifications	Ubiquitinated; mediated by SIAH1 or SIAH2 and leading to its subsequent proteasomal degradation.
Cellular localization	Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane. Cell junction, synapse, synaptosome.

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab267272 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 33 kDa.

Images



All lanes : Anti-Synaptophysin antibody [EP1098Y] ([ab52636](#)) at 1/1000 dilution

Lane 1 : Wild-type HEK293T cell lysate

Lane 2 : Synp knockout HEK293T cell lysate

Lane 3 : Human brain tissue lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 33 kDa

Observed band size: 38 kDa

Lanes 1-3: Merged signal (red and green). Green - [ab52636](#) observed at 38 kDa. Red - loading control, [ab7291](#) observed at 50 kDa.

[ab52636](#) Anti-Synaptophysin antibody [EP1098Y] was shown to specifically react with Synaptophysin in wild-type HEK293T cells. Loss of signal was observed when knockout cell line ab267272 (knockout cell lysate [ab257060](#)) was used. Wild-type and Synaptophysin knockout samples were subjected to SDS-PAGE. [ab52636](#) and Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti- Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti- Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.



Western blot - Human Syp knockout HEK293T cell line (ab267272)

All lanes : Anti-Synaptophysin antibody [YE269] (**ab32127**) at 1/1000 dilution

Lane 1 : Wild-type HEK293T cell lysate

Lane 2 : Syp knockout HEK293T cell lysate

Lane 3 : Human brain tissue lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 33 kDa

Observed band size: 38 kDa


Lanes 1-3: Merged signal (red and green). Green - **ab32127** observed at 38 kDa. Red - loading control, **ab7291** observed at 50 kDa.

ab32127 Anti-Synaptophysin antibody [YE269] was shown to specifically react with Synaptophysin in wild-type HEK293T cells. Loss of signal was observed when knockout cell line ab267272 (knockout cell lysate **ab257060**) was used. Wild-type and Synaptophysin knockout samples were subjected to SDS-PAGE. **ab32127** and Anti-alpha Tubulin antibody [DM1A] - Loading Control (**ab7291**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti- Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti- Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.



Sanger Sequencing - Human Syp knockout HEK293T cell line (ab267272)

Homozygous: 8 bp deletion in exon2



Representative images of SYP knockout HEK293T cells, low and high confluency examples (top left and right respectively) and wild-type HEK293T cells, low and high confluency (bottom left and right respectively) showing typical adherent, epithelial-like morphology. Images were captured at 10X magnification using a EVOS XL Core microscope.

Cell Culture - Human SYP (Synaptophysin)
knockout HEK293T cell line (ab267272)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors