

# Human UCHL1 (PGP9.5) knockout HEK-293T cell line ab255443

2 Images

### Overview

<b>Product name</b>	Human UCHL1 (PGP9.5) knockout HEK-293T cell line
<b>Parental Cell Line</b>	HEK293T
<b>Organism</b>	Human
<b>Mutation description</b>	Knockout achieved by using CRISPR/Cas9, Homozygous: 45 bp deletion in exon 1
<b>Passage number</b>	<20
<b>Knockout validation</b>	Sanger Sequencing, Western Blot (WB)
<b>Tested applications</b>	<b>Suitable for:</b> WB
<b>Biosafety level</b>	2
<b>General notes</b>	<p><b>Recommended control:</b> Human wild-type HEK293T cell line (<a href="#">ab255593</a>). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p><b>Cryopreservation cell medium:</b> Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p><b>Culture medium:</b> DMEM (High Glucose) + 10% FBS</p> <p><b>Initial handling guidelines:</b> Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none"> <li>1. Thaw the vial in 37°C water bath for approximately 1-2 minutes.</li> <li>2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.</li> <li>3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of <math>2 \times 10^4</math> cells/cm<sup>2</sup>. Seeding density is given as a guide only and should be scaled to align with individual lab schedules.</li> <li>4. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>. Cultures should be monitored daily.</li> </ol> <p><b>Subculture guidelines:</b></p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of <math>2 \times 10^4</math> cells/cm<sup>2</sup> is recommended.</p>

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.

Cells should be passaged when they have achieved 80-90% confluence.

This product is subject to limited use licenses from The Broad Institute and ERS Genomics Limited, and is developed with patented technology. For full details of the limited use licenses and relevant patents please refer to our [limited use license](#) and [patent pages](#).

We will provide viable cells that proliferate on revival.

## Properties

---

<b>Number of cells</b>	1 x 10 <sup>6</sup> cells/vial, 1 mL
<b>Adherent /Suspension</b>	Adherent
<b>Tissue</b>	Kidney
<b>Cell type</b>	epithelial
<b>STR Analysis</b>	Amelogenin X D5S818: 8, 9 D13S317: 11, 12, 14 D7S820: 11 D16S539: 9, 13 vWA: 15, 20 TH01: 7, 9.3 TPOX: 11, 12 CSF1PO: 12
<b>Mycoplasma free</b>	Yes
<b>Storage instructions</b>	Shipped on Dry Ice. Store in liquid nitrogen.
<b>Storage buffer</b>	Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether

## Target

---

<b>Function</b>	Ubiquitin-protein hydrolase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. Also binds to free monoubiquitin and may prevent its degradation in lysosomes. The homodimer may have ATP-independent ubiquitin ligase activity.
<b>Tissue specificity</b>	Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.
<b>Involvement in disease</b>	Parkinson disease 5 Neurodegeneration with optic atrophy, childhood-onset
<b>Sequence similarities</b>	Belongs to the peptidase C12 family.
<b>Post-translational modifications</b>	O-glycosylated.
<b>Cellular localization</b>	Cytoplasm. Endoplasmic reticulum membrane. About 30% of total UCHL1 is associated with membranes in brain.


## Applications

---

**The Abpromise guarantee** Our [Abpromise guarantee](#) covers the use of ab255443 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 24 kDa.

## Images



**All lanes :** Anti-PGP9.5 antibody [EPR4118] - Neuronal Marker ([ab108986](#)) at 1/1000 dilution

**Lane 1 :** Wild-type HAP1 cell lysate

**Lane 2 :** UCHL1 knockout HAP1 cell lysate

**Lane 3 :** SH-SY5Y cell lysate

**Lane 4 :** Wild-type HEK-293T cell lysate

**Lane 5 :** UCHL1 knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/20000 dilution

Performed under reducing conditions.

**Predicted band size:** 24 kDa

**Additional bands at:** 37 kDa (possible Loading Control)

**Lanes 1 - 5:** Merged signal (red and green). Green - [ab108986](#) observed at 25 kDa. Red - loading control, [ab8245](#) observed at 37 kDa.

[ab108986](#) was shown to react with PGP9.5 in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line ab255443 (knockout cell lysate [ab263773](#)) was used. Wild-type and PGP9.5 knockout samples were subjected to SDS-PAGE. [ab108986](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4 °C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

Mut	.....GGTGCACCGCTACCC
WT	AGATGCAGCTCAAGCCGATGGAGATCAACCCGAGGTGAGCGCCAGGTGCACCGCTACCC

Sanger Sequencing - Human UCHL1 knockout  
HEK293T cell line (ab255443)

Homozygous: 45 bp deletion in exon 1.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors