abcam

Product datasheet

Human BIRC2 (cIAP1) knockout HeLa cell lysate ab257372

2 Images

Overview

Product name	Human BIRC2 (cIAP1) knockout HeLa cell lysate
Product overview	Knockout cell lysate achieved by CRISPR/Cas9.
Parental Cell Line	HeLa
Organism	Human
Mutation description	Knockout achieved by using CRISPR/Cas9, 1 bp deletion in exon2.
Passage number	<20
Knockout validation	Sanger Sequencing, Western Blot (WB)
Reconstitution notes	To use as WB control, resuspend the lyophilizate in 50 µL of LDS* Sample Buffer to have a final concentration of 2 mg/ml. For reducing conditions, we recommend a final concentration of 0.1 M DTT. **Usage of SDS sample buffer is not recommended with these lyophilized lysates.*
Notes	**Lysate preparation:** Our lysates are made using RIPA buffer to which we add a protease inhibitor cocktail and phosphatase inhibitor cocktail (ratio: 300:100:10). *This means that the protein of interest is denatured.* If you require a native form of the protein please use the live cell version - found **here**. Please refer to our lysis protocol for further details on how our lysates are prepared. **User storage instructions:** Lyophilizate may be stored at 4°C. After reconstitution, store at -20°C for short-term storage or -80°C for long-term storage. Access thousands of knockout cell lysates, generated from commonly used cancer cell lines. **See here for more information on knockout cell lysates.** Abcam has not and does not intend to apply for the REACH Authorisation of customers' uses of products that contain European Authorisation list (Annex XIV) substances. It is the responsibility of our customers to check the necessity of application of REACH Authorisation, and any other relevant authorisations, for their intended uses. This product is subject to limited use licenses from The Broad Institute and ERS Genomics Limited, and is developed with patented technology. For full details of the limited use licenses and relevant patents please refer to our **limited use license** and **patent pages**.
Tested applications	**Suitable for:** WB

Properties

Storage instructions Store at -80°C. Please refer to protocols.

Components	1 kit
ab260212 - Human BIRC2 knockout HeLa cell lysate	1 x 100µg
ab255929 - Human wild-type HeLa cell lysate	1 x 100µg

Cell type epithelial

Disease Adenocarcinoma

Gender Female

STR Analysis Amelogenin X D5S818: 11, 12 D13S317: 12, 13.3 D7S820: 8, 12 D16S539: 9, 10 vWA: 16, 18 TH01: 7 TPOX: 8,12 CSF1PO: 9, 10

Target

Function Apoptotic suppressor. The BIR motifs region interacts with TNF receptor associated factors 1 and 2 (TRAF1 and TRAF2) to form an heteromeric complex, which is then recruited to the tumor necrosis factor receptor 2 (TNFR2).

Tissue specificity Present in many fetal and adult tissues. Mainly expressed in adult skeletal muscle, thymus, testis, ovary, and pancreas, low or absent in brain and peripheral blood leukocytes.

Sequence similarities Belongs to the IAP family.
Contains 3 BIR repeats.
Contains 1 CARD domain.
Contains 1 RING-type zinc finger.

Cellular localization Cytoplasm.

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab257372 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 70 kDa.

Images



Western blot - Human BIRC2 (cIAP1) knockout HeLa cell lysate (ab257372)

Lane 1: Wild-type HeLa cell lysate (20µg)

Lane 2: BIRC2 knockout HeLa cell lysate (20µg)

Lanes 1- 2: Merged signal (red and green). Green - **ab108361** observed at 70 kDa. Red - loading control **ab8245** observed at 37 kDa.

ab108361 Anti-cIAP1 antibody [EPR4673] was shown to specifically react with cIAP1 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line **ab265896** (knockout cell lysate ab257372) was used. Wild-type and cIAP1 knockout samples were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. **ab108361** and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 1000 and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

Sanger Sequencing - Human BIRC2 knockout HeLa cell lysate (ab257372)

Homozygous: 1 bp deletion in exon2

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit **https://www.abcam.com/abpromise** or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors