abcam

Product datasheet

SW 1353 whole cell lysate ab7915

1 References

Overview

Product name SW 1353 whole cell lysate

General notes Cell line: SW1353 (Human humerus chondrosarcoma, fibroblast-like).

Growth media: DMEM and 10% FBS (Fetal bovine serum).

SW1353 cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium

chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM

phenylmethylsulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium

dodecylsulfate, 5 μ g/ml of aprotinin, 5 μ g/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate

was boiled for 5 min in 1 x SDS sample buffer (0.045 M Tris-HCl pH 6.8, 10% glycerol, 1%

sodium dodecylsulfate, 0.01% bromophenol blue), containing 0.05 M DTT.

Tested applications Suitable for: WB

Properties

Mycoplasma free Yes

Form Liquid

Storage instructions Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.

Storage buffer pH: 7.20

Constituent: 100% SDS Sample Buffer

Lysate notes SW1353 cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium

chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM $\,$

phenylmethylsulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by

centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1%

sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.

Background The SW1353 chondrosarcoma cell line is considered to be a model for studies of the

transcription factor Sox9, an important determinant of chondrocyte differentiation, including the regulation of type II collagen (Col2) and aggrecan gene expression. Cells are used to study

mechanisms of joint inflammation and cartilage destruction.

1

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab7915 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. SW1353 cell lysate is ready to load on SDS-PAGE for Western blotting, 20 µg per lane is recommended for mini gel.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- · Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors