abcam

Product datasheet

Mouse AGT Antibody Pair - BSA and Azide free ab256659



2 Images

Overview

Product name Mouse AGT Antibody Pair - BSA and Azide free

Assay type ELISA set

0.057 pg/ml - 10000 pg/ml Range

Species reactivity Reacts with: Mouse

Product overview The Antibody Pair can be used to quantify Mouse AGT. BSA and Azide free antibody pairs

> include unconjugated capture and detector antibodies suitable for sandwich ELISAs. The antibodies are provided at an approximate concentration of 1 mg/ml as measured by the protein A280 method. The recommended antibody orientation is based on internal optimization for ELISA-based assays. Antibody orientation is assay dependent and needs to be optimized for each assay type. Both capture and detector antibodies are rabbit monoclonal antibodies

delivering consistent, specific, and sensitive results.

For additional information on the performance of the antibody pair, see the equivalent SimpleStep ELISA® Kit (ab245718), which uses the same antibodies. However, due to differences in their formulation, this antibody pair cannot be used with the consumables provided with our SimpleStep ELISA Kits. Please note that the range provided for the pairs is only an estimation based on the performance of the related product using the same antibody pair. Performance of the antibody pair will depend on the specific characteristics of your assay. We guarantee the product works in sandwich ELISA, but we do not guarantee the sensitivity or dynamic range of the antibody pair in your assay.

Download SDS here.

Tested applications Suitable for: Sandwich ELISA

Platform Reagents

Properties

Storage instructions Store at +4°C. Please refer to protocols.

Carrier free Yes

Components	10 x 96 tests
Mouse AGT Capture Antibody (unconjugated)	1 x 100µg
Mouse AGT Detector Antibody (unconjugated)	1 x 100µg

Function

Essential component of the renin-angiotensin system (RAS), a potent regulator of blood pressure, body fluid and electrolyte homeostasis. In response to lowered blood pressure, the enzyme renin cleaves angiotensinogen to produce angiotensin-1 (angiotensin 1-10). Angiotensin-1 is a substrate of ACE (angiotensin converting enzyme) that removes a dipeptide to yield the physiologically active peptide angiotensin-2 (angiotensin 1-8). Angiotensin-1 and angiotensin-2 can be further processed to generate angiotensin-3 (angiotensin 2-8), angiotensin-4 (angiotensin 3-8). Angiotensin 1-7 is cleaved from angiotensin-2 by ACE2 or from angiotensin-1 by MME (neprilysin). Angiotensin 1-9 is cleaved from angiotensin-1 by ACE2.

Angiotensin-2 acts directly on vascular smooth muscle as a potent vasoconstrictor, affects cardiac contractility and heart rate through its action on the sympathetic nervous system, and alters renal sodium and water absorption through its ability to stimulate the zona glomerulosa cells of the adrenal cortex to synthesize and secrete aldosterone.

Angiotensin-3 stimulates aldosterone release.

Angiotensin 1-7 is a ligand for the G-protein coupled receptor MAS1 (By similarity). Has vasodilator and antidiuretic effects (By similarity). Has an antithrombotic effect that involves MAS1-mediated release of nitric oxide from platelets.

Tissue specificity

Expressed by the liver and secreted in plasma.

Involvement in disease

Genetic variations in AGT are a cause of susceptibility to essential hypertension (EHT)

[MIM:145500]. Essential hypertension is a condition in which blood pressure is consistently higher

than normal with no identifiable cause.

Defects in AGT are a cause of renal tubular dysgenesis (RTD) [MIM:267430]. RTD is an autosomal recessive severe disorder of renal tubular development characterized by persistent fetal anuria and perinatal death, probably due to pulmonary hypoplasia from early-onset

oligohydramnios (the Potter phenotype).

Sequence similarities

Belongs to the serpin family.

Post-translational modifications

Beta-decarboxylation of Asp-34 in angiotensin-2, by mononuclear leukocytes produces alanine.

The resulting peptide form, angiotensin-A, has the same affinity for the AT1 receptor as

angiotensin-2, but a higher affinity for the AT2 receptor.

Cellular localization Secreted.

Applications


The Abpromise guarantee

Our Abpromise guarantee covers the use of ab256659 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
Sandwich ELISA		Use at an assay dependent concentration.

Images



Sandwich ELISA - Mouse AGT Antibody Pair - BSA and Azide free (ab256659)

Representative standard curve from corresponding SimpleStep ELISA® Kit (ab245718), which uses the same antibody pair. For additional information on the performance of pair and kit, refer to the corresponding kit datasheet. Due to differences in the formulation and format of the antibodies in this pair, they cannot be used as substitutes for the antibody components in our SimpleStep ELISA® Kits.



To learn more about the advantages of recombinant antibodies see **here**.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors