

Product datasheet

Anti-ABCD1/ALD antibody [EPR15929] ab197013

Recombinant RabMAb

★★★★☆ [1 Abreviews](#) [4 References](#) [6 Images](#)

Overview

| | |
|----------------------------|--|
| Product name | Anti-ABCD1/ALD antibody [EPR15929] |
| Description | Rabbit monoclonal [EPR15929] to ABCD1/ALD |
| Host species | Rabbit |
| Tested applications | Suitable for: WB, ICC/IF, Flow Cyt (Intra) |
| Species reactivity | Reacts with: Mouse, Rat, Human |
| Immunogen | Recombinant fragment. This information is proprietary to Abcam and/or its suppliers. |
| Positive control | WB: HepG2 cells, HeLa cells, human fetal liver lysate; JAR cells; C6 cells, RAW 264.7 cells, PC-12 cells, NIH/3t3 cells. ICC/IF: MCF7 cells. Flow Cyt (intra): HeLa cells. |
| General notes | <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p> |

Properties

| | |
|-----------------------------|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. |
| Storage buffer | pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA |

| | |
|---------------------|--------------------|
| Purity | Protein A purified |
| Clonality | Monoclonal |
| Clone number | EPR15929 |
| Isotype | IgG |

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab197013 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------------------|-----------|---|
| WB | | 1/10000. Detects a band of approximately 83 kDa (predicted molecular weight: 83 kDa). |
| ICC/IF | | 1/100. |
| Flow Cyt (Intra) | | Use at an assay dependent concentration. |

Target

| | |
|-------------------------------|---|
| Function | Probable transporter. The nucleotide-binding fold acts as an ATP-binding subunit with ATPase activity. |
| Involvement in disease | Defects in ABCD1 are the cause of adrenoleukodystrophy X-linked (X-ALD) [MIM:300100]. X-ALD is a peroxisomal metabolic disorder characterized by progressive multifocal demyelination of the central nervous system and by peripheral adrenal insufficiency (Addison disease). It results in mental deterioration, corticospinal tract dysfunction, and cortical blindness. Different clinical manifestations exist like: cerebral childhood ALD (CALD), adult cerebral ALD (ACALD), adrenomyeloneuropathy (AMN) and 'Addison disease only' (ADO) phenotype. Note=The promoter region of ABCD1 is deleted in the chromosome Xq28 deletion syndrome which involves ABCD1 and the neighboring gene BCAP31. |
| Sequence similarities | Belongs to the ABC transporter superfamily. ABCD family. Peroxisomal fatty acyl CoA transporter (TC 3.A.1.203) subfamily. Contains 1 ABC transmembrane type-1 domain. Contains 1 ABC transporter domain. |
| Cellular localization | Peroxisome membrane. |

Images



All lanes : Anti-ABCD1/ALD antibody [EPR15929] (ab197013) at 1/10000 dilution

Lane 1 : HepG2 (Human liver hepatocellular carcinoma) whole cell lysate

Lane 2 : HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysate

Lane 3 : Human fetal liver lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 83 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized MCF7 (Human breast adenocarcinoma cell line) cells labeling ABCD1 / ALD with ab197013 at 1/100 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/500 dilution (green). Cytoplasm staining on MCF7 cell line is observed. The nuclear counter stain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:

-ve control 1: ab197013 at 1/100 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.

-ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.




Flow Cytometry (Intracellular) - Anti-ABCD1/ALD antibody [EPR15929] (ab197013)

ab197013 staining ABCD1/ALD in the human cell line HeLa (human cervix adenocarcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. The sample was incubated with the primary antibody at a dilution of 1/70. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isotype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)



Western blot - Anti-ABCD1/ALD antibody [EPR15929] (ab197013)

Anti-ABCD1/ALD antibody [EPR15929] (ab197013) at 1/10000 dilution + JAR (Human placenta choriocarcinoma cell line) whole cell lysate at 20 µg


Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 83 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-ABCD1/ALD antibody [EPR15929] (ab197013)

All lanes : Anti-ABCD1/ALD antibody [EPR15929] (ab197013) at 1/10000 dilution

Lane 1 : C6 (Rat glial tumor cells) whole cell lysate

Lane 2 : RAW 264.7 (Mouse macrophage cells transformed with Abelson murine leukemia virus) whole cell lysate

Lane 3 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate

Lane 4 : NIH/3T3 (Mouse embryo fibroblast cells) whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary



All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 83 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.

Why choose a recombinant antibody?

| | |
|---|---|
|  Research with confidence Consistent and reproducible results |  Long-term and scalable supply Recombinant technology |
|  Success from the first experiment Confirmed specificity |  Ethical standards compliant Animal-free production |

Anti-ABCD1/ALD antibody [EPR15929] (ab197013)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors