


Product datasheet

Anti-Abhd5/CGI-58 antibody ab111984

[1 References](#) [3 Images](#)

Overview

Product name	Anti-Abhd5/CGI-58 antibody
Description	Goat polyclonal to Abhd5/CGI-58
Host species	Goat
Tested applications	Suitable for: WB, IHC-P
Species reactivity	Reacts with: Mouse, Human Predicted to work with: Rat, Dog, Pig 
Immunogen	Synthetic peptide corresponding to Human Abhd5/CGI-58 aa 150-250 (internal sequence). Database link: Q8WTS1

 [Run BLAST with](#)

 [Run BLAST with](#)

General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
Storage buffer	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 0.5% BSA, 99% Tris buffered saline
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab111984 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use a concentration of 0.2 - 0.6 µg/ml. Predicted molecular weight: 39 kDa.
IHC-P		Use a concentration of 3.75 µg/ml.

Target**Function**

Lysophosphatidic acid acyltransferase which functions in phosphatidic acid biosynthesis. May regulate the cellular storage of triacylglycerol through activation of the phospholipase PNPLA2. Involved in keratinocyte differentiation.

Tissue specificity

Widely expressed in various tissues, including lymphocytes, liver, skeletal muscle and brain. Expressed by upper epidermal layers and dermal fibroblasts in skin, hepatocytes and neurons (at protein level).

Involvement in disease

Defects in ABHD5 are the cause of Chanarin-Dorfman syndrome (CDS) [MIM:275630]; also called triglyceride storage disease with impaired long-chain fatty acid oxidation or neutral lipid storage disease with ichthyosis. CDS is an autosomal recessive inborn error of lipid metabolism with multisystemic accumulation of triglycerides although plasma concentrations are normal. Clinical characteristics are congenital generalized ichthyosis, vacuolated leukocytes, hepatomegaly, myopathy, cataracts, neurosensory hearing loss and developmental delay. The disorder presents at birth with generalized, fine, white scaling of the skin and a variable degree of erythema resembling non-bullous congenital ichthyosiform erythroderma.

Sequence similarities

Belongs to the peptidase S33 family. ABHD4/ABHD5 subfamily.

Developmental stage

Detected in fetal epidermis from 49 to 135 days estimated gestational age (at protein level).


Domain

The HXXXXD motif is essential for acyltransferase activity and may constitute the binding site for the phosphate moiety of the glycerol-3-phosphate.

Cellular localization


Cytoplasm. Lipid droplet. Colocalized with PLIN and ADRP on the surface of lipid droplets. The localization is dependent upon the metabolic status of the adipocytes and the activity of PKA.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Abhd5/CGI-58 antibody (ab111984)

ab111984, at 3.75 µg/ml, staining Abhd5/CGI-58 in formalin-fixed, paraffin-embedded Human kidney tissue by Immunohistochemistry.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Abhd5/CGI-58 antibody (ab111984)

ab111984, at 3.75 µg/ml, staining Abhd5/CGI-58 in formalin-fixed, paraffin-embedded Human pancreas tissue by Immunohistochemistry.



Western blot - Anti-Abhd5/CGI-58 antibody (ab111984)

Anti-Abhd5/CGI-58 antibody (ab111984) at 0.2 µg/ml + NIH3T3 lysate (in RIPA buffer) at 35 µg

Developed using the ECL technique.

Predicted band size: 39 kDa

Primary incubation was 1 hour.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors