

## Product datasheet

# Anti-Ada protein antibody [ADA-1] ab18104

## 2 References

### Overview

---

<b>Product name</b>	Anti-Ada protein antibody [ADA-1]
<b>Description</b>	Mouse monoclonal [ADA-1] to Ada protein
<b>Host species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> RIA, WB, IP
<b>Species reactivity</b>	<b>Reacts with:</b> Escherichia coli
<b>Immunogen</b>	Full length protein corresponding to Mouse Ada protein. Please refer to the reference below.
<b>General notes</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Preservative: 0.02% Sodium azide Constituent: 99.98% PBS
<b>Purity</b>	Protein A/G purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	ADA-1
<b>Isotype</b>	unknown

### Applications

---

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab18104 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
RIA		Use at an assay dependent dilution.
WB		Use at an assay dependent dilution. Predicted molecular weight: 40 kDa.
IP		Use at an assay dependent dilution.

## Target

---

### Relevance

The methylated ADA protein acts as a positive regulator of its own synthesis, as well as that of other proteins. The transcription-activating function of the ADA protein resides in its N-terminus. It activates the transcription of *alkA*, *alkB* and *aidB*.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors