# abcam

### Product datasheet

## Anti-ADRA1B antibody ab84405

lgG

★★★★★ <u>1 Abreviews</u> <u>3 References</u> 3 Images

#### Overview

| Product name         | Anti-ADRA1B antibody  |  |  |
|----------------------|---|--|--|
| Description          | Rabbit polyclonal to ADRA1B   |  |  |
| Host species         | Rabbit  |  |  |
| Tested applications  | Suitable for: ICC/IF, WB, IHC-P   |  |  |
| Species reactivity   | Reacts with: Human  |  |  |
| Immunogen            | Synthetic peptide corresponding to Human ADRA1B (C terminal). (NM_000679; P35368)<br>(Peptide available as <u>ab226906</u> )  |  |  |
| Positive control     | Human fetal brain lysates; human heart tissue.  |  |  |
| General notes        | The Life Science industry has been in the grips of a reproducibility crisis for a number of years.<br>Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies<br>and knockout edited cell lines for gold-standard validation. Please check that this product meets<br>your needs before purchasing. |  |  |
|                      | If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As  |  |  |
| Properties           |   |  |  |
| Form                 | Lyophilized:Reconstitute with 200ul distilled sterile water. Please note that if you receive this product in liquid form it has already been reconstituted as described and no further reconstitution is necessary.   |  |  |
| Storage instructions | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.  |  |  |
| Storage buffer       | Preservative: 0.02% Sodium azide<br>Constituent: 1% BSA   |  |  |
| Purity               | Immunogen affinity purified   |  |  |
| Purification notes   | ab84405 is purified by a peptide affinity column.   |  |  |
| Clonality            | Polyclonal  |  |  |

lsotype

#### The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab84405 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews            | Notes   |
|-------------|----------------------|---|
| ICC/IF      |                      | Use a concentration of 1 µg/ml.   |
| WB          |                      | 1/200 - 1/1000. Predicted molecular weight: 57 kDa.Can be blocked with <b>Human ADRA1B peptide (ab226906)</b> . |
| IHC-P       | <b>★ ★ ★ ★ ☆ (1)</b> | 1/100 - 1/500.  |


| Target                |  |
|-----------------------|--|
| Function              | This alpha-adrenergic receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. Its effect is mediated by G(q) and G(11) proteins. Nuclear ADRA1A-ADRA1B heterooligomers regulate phenylephrine (PE)-stimulated ERK signaling in cardiac myocytes. |
| Sequence similarities | Belongs to the G-protein coupled receptor 1 family. Adrenergic receptor subfamily. ADRA1B sub-<br>subfamily.   |
| Cellular localization | Nucleus membrane. Cell membrane. Location at the nuclear membrane facilitates<br>heterooligomerization and regulates ERK-mediated signaling in cardiac myocytes. signaling in<br>cardiac myocytes. Colocalizes with GNAQ, PLCB1 as well as LAP2 at the nuclear membrane of<br>cardiac myocytes.                            |

#### Images

. .




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ADRA1B antibody (ab84405) ab84405, at a 1/100 dilution, staining ADRA1B in formalin fixed, paraffin embedded human heart tissue by Immunohistochemistry.



Anti-ADRA1B antibody (ab84405) at 1/500 dilution + human fetal brain lysates

Predicted band size: 57 kDa Observed band size: 57 kDa

Western blot - Anti-ADRA1B antibody (ab84405)



Immunocytochemistry/ Immunofluorescence - Anti-ADRA1B antibody (ab84405) ICC/IF image of <u>ab84450</u> stained HeLa cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (<u>ab84450</u>, 1µg/ml) overnight at +4°C. The secondary antibody (green) was <u>ab96899</u>, DyLight® 488 goat anti-rabbit lgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors