abcam

Product datasheet

Alexa Fluor® 555 Anti-LAG-3 antibody [EPR20261] ab280858

Recombinant

2 Images

Overview

Product name Alexa Fluor® 555 Anti-LAG-3 antibody [EPR20261]

Description Alexa Fluor® 555 Rabbit monoclonal [EPR20261] to LAG-3

Host species Rabbit

Conjugation Alexa Fluor® 555. Ex: 555nm, Em: 565nm

Tested applications

Suitable for: IHC-P

Species reactivity

Reacts with: Human

Immunogen Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

Positive control IHC-P: Human Hodgkin's lymphoma tissue.

General notes This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor[®] dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or **outlicensing@thermofisher.com**.

1

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle. Store In the Dark.

Storage buffer pH: 7.4

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, 68.98% PBS

Purity Protein A purified

ClonalityMonoclonalClone numberEPR20261

Isotype IgG

Applications

The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab280858 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		1/50. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target

Function Involved in lymphocyte activation. Binds to HLA class-II antigens.


Tissue specificity On cell surface of activated NK and T-lymphocytes.

Sequence similarities Contains 3 lg-like C2-type (immunoglobulin-like) domains.

Contains 1 lg-like V-type (immunoglobulin-like) domain.

Cellular localization Membrane.

Images



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Alexa Fluor® 555 Anti-LAG-3 antibody [EPR20261] (ab280858)


Immunofluorescence staining of LAG-3 in a section of formalinfixed, paraffin-embedded human Hodgkin's lymphoma*.

Performed on a Leica BOND[®]. The section was pre-treated using heat mediated antigen retrieval with EDTA (pH 9.0) using retrieval settings of 110°C for 20 minutes. The section was then incubated at room temperature for 1 hour with ab280858 at 1/50 dilution (shown in green) and counterstained using **ab190573**, Rabbit anti-Tubulin (A647)(1/250) at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue). The section was then mounted using Dako Fluorescence Mounting Medium [®].

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

For other IHC staining systems (automated and non-automated), customers should optimize variable parameters such as antigen retrieval conditions, antibody concentrations and incubation times.

*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.



(ab280858)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors