abcam

Product datasheet

Alexa Fluor® 555 Anti-STAT3 (phospho Y705) antibody [EP2147Y] ab202876

Recombinant RabMAb

★★★★★ 1 Abreviews 1 References 2 Images

Overview

Product name Alexa Fluor® 555 Anti-STAT3 (phospho Y705) antibody [EP2147Y]

Alexa Fluor® 555 Rabbit monoclonal [EP2147Y] to STAT3 (phospho Y705) **Description**

Host species Rabbit

Conjugation Alexa Fluor® 555. Ex: 555nm, Em: 565nm

Tested applications Suitable for: ICC/IF Species reactivity Reacts with: Human

Predicted to work with: Mouse, Rat

Immunogen Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

(Peptide available as ab179551)

Positive control ICC/IF: HeLa cells treated with 50ng/ml IFNa for 5 minutes

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit General notes

monoclonal antibodies. For details on our patents, please refer to **RabMAb patents**.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Stable for 12 months at -20°C. Store In the Dark.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA

Purity Protein A purified

Clonality Monoclonal
Clone number EP2147Y

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab202876 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF	★ ★ ★ ★ (1)	1/100. This product gave a positive signal in HeLa cells treated with 50ng/ml IFN? for 5 minutes and fixed with 4% formaldehyde (10 min).

Target

Function

Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors. Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (PubMed:17344214). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the interleukin-6 (IL-6)-responsive elements identified in the promoters of various acute-phase protein genes. Activated by IL31 through IL31RA. Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed:17344214). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation (By similarity). May play an apoptotic role by transctivating BIRC5 expression under LEP activation (PubMed:18242580). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity.

Tissue specificity

Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.

Involvement in disease

Hyperimmunoglobulin E recurrent infection syndrome, autosomal dominant

Autoimmune disease, multisystem, infantile-onset

Sequence similarities

Belongs to the transcription factor STAT family.

Contains 1 SH2 domain.

Post-translational modifications

Tyrosine phosphorylated upon stimulation with EGF. Tyrosine phosphorylated in response to constitutively activated FGFR1, FGFR2, FGFR3 and FGFR4 (By similarity). Activated through tyrosine phosphorylation by BMX. Tyrosine phosphorylated in response to IL6, IL11, LIF, CNTF, KITLG/SCF, CSF1, EGF, PDGF, IFN-alpha, LEP and OSM. Activated KIT promotes


phosphorylation on tyrosine residues and subsequent translocation to the nucleus.

Phosphorylated on serine upon DNA damage, probably by ATM or ATR. Serine phosphorylation is important for the formation of stable DNA-binding STAT3 homodimers and maximal transcriptional activity. ARL2BP may participate in keeping the phosphorylated state of STAT3 within the nucleus. Upon LPS challenge, phosphorylated within the nucleus by IRAK1. Upon erythropoietin treatment, phosphorylated on Ser-727 by RPS6KA5. Phosphorylation at Tyr-705 by PTK6 or FER leads to an increase of its transcriptional activity. Dephosphorylation on tyrosine residues by PTPN2 negatively regulates IL6/interleukin-6 signaling.

Cellular localization


Cytoplasm. Nucleus. Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4. Constitutive nuclear presence is independent of tyrosine phosphorylation. Predominantly present in the cytoplasm without stimuli. Upon leukemia inhibitory factor (LIF) stimulation, accumulates in the nucleus. The complex composed of BART and ARL2 plays an important role in the nuclear translocation and retention of STAT3. Identified in a complex with LYN and PAG1.

Images



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 555 Anti-STAT3 (phospho Y705) antibody [EP2147Y] (ab202876)

ab202876 staining STAT3 (phospho Y705) in HeLa cells. The cells were incubated with 50ng/ml IFNα for 5 minutes (Treated) or solvent-only for control purposes (Non-treated). Cells were fixed with 4% formaldehyde (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab202877 at 1/100 dilution (pseudocolored in red) and ab195887 Mouse monoclonal [DM1A] to alpha Tubulin - Microtubule Marker (Alexa Fluor® 488) at 2μg/ml (shown in green) overnight at +4°C. Nuclear DNA was labelled in blue with DAPI. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Alexa Fluor® 555 Anti-STAT3 (phospho Y705) antibody [EP2147Y] (ab202876)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors