abcam

Product datasheet

Alexa Fluor® 647 Anti-Cytokeratin 18 antibody [EPR1626] ab206269

Recombinant RabMAb

2 Images

Overview

Product name Alexa Fluor® 647 Anti-Cytokeratin 18 antibody [EPR1626]

Description Alexa Fluor® 647 Rabbit monoclonal [EPR1626] to Cytokeratin 18

Host species Rabbit

Conjugation Alexa Fluor® 647. Ex: 652nm. Em: 668nm

Tested applications Suitable for: ICC/IF Species reactivity Reacts with: Human

Predicted to work with: Rat ...

Synthetic peptide corresponding to Human Cytokeratin 18 (C terminal). **Immunogen**

Database link: P05783

Positive control ICC/IF: HepG2 cells

General notes This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

Purity Protein A purified

Clonality Monoclonal Clone number **EPR1626**

Isotype lqG

Applications

Target

The Abpromise guarantee

Our Abpromise guarantee covers the use of ab206269 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/100. This product gave a positive signal in HepG2 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

Function Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (E
--

phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier

protection.

Tissue specificity Expressed in colon, placenta, liver and very weakly in exocervix. Increased expression observed

in lymph nodes of breast carcinoma.

Involvement in disease Defects in KRT18 are a cause of cirrhosis (CIRRH) [MIM:215600].

Sequence similarities Belongs to the intermediate filament family.

Post-translational Phosphorylation at Ser-34 increases during mitosis. Hyperphosphorylated at Ser-53 in diseased modifications


cirrhosis liver. Phosphorylation increases by IL-6.

Proteolytically cleaved by caspases during epithelial cell apoptosis. Cleavage occurs at Asp-238

by either caspase-3, caspase-6 or caspase-7.

O-glycosylated at multiple sites; glycans consist of single N-acetylglucosamine residues.


Cellular localization Cytoplasm > perinuclear region.



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-Cytokeratin 18 antibody [EPR1626] (ab206269) ab206269 staining Cytokeratin 18 in HepG2 cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab206269 at a 1/100 dilution (shown in red) and ab195887, Mouse monoclonal to alpha Tubulin (Alexa Fluor[®] 488), at a 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HepG2 cells fixed with 4% formaldehyde (10 min).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors