# abcam

## Product datasheet

# Alexa Fluor® 647 Anti-MCM2 antibody [EPR4120] ab223403



#### 1 References 3 Images

#### Overview

**Product name** Alexa Fluor® 647 Anti-MCM2 antibody [EPR4120]

**Description** Alexa Fluor® 647 Rabbit monoclonal [EPR4120] to MCM2

**Host species** Rabbit

Alexa Fluor® 647. Ex: 652nm, Em: 668nm Conjugation

**Tested applications** Suitable for: ICC/IF, Flow Cyt (Intra)

Species reactivity Reacts with: Human

Predicted to work with: Mouse, Rat

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. **Immunogen** 

Positive control ICC/IF: MCF7 cells. Flow Cyt (intra): MCF7 cells

**General notes** This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor<sup>®</sup> is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor<sup>®</sup> dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

1

#### outlicensing@thermofisher.com.

#### **Properties**

**Form** Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 1% BSA, 30% Glycerol (glycerin, glycerine), PBS

**Purity** Protein A purified

Clonality Monoclonal Clone number EPR4120

Isotype lgG

#### **Applications**

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab223403 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/100. This product gave a positive signal in MCF7 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min)
Flow Cyt (Intra)		1/5000.

### **Target**

**Function** Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative

> helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase

activity. Required for the entry in S phase and for cell division.

Sequence similarities Belongs to the MCM family.

Contains 1 MCM domain.

Post-translational


Phosphorylated on Ser-108 by ATR in proliferating cells. Ser-108 proliferation is increased by modifications

genotoxic agents. Ser-40 is mediated by the CDC7-DBF4 and CDC7-DBF4B complexes, while Ser-53 phosphorylation is only mediated by the CDC7-DBF4 complex. Phosphorylation by the

CDC7-DBF4 complex during G1/S phase is required for the initiation of DNA replication.

**Cellular localization** Nucleus.

#### **Images**




Flow Cytometry (Intracellular) - Alexa Fluor® 647 Anti-MCM2 antibody [EPR4120] (ab223403)

Overlay histogram showing MCF7 cells stained with ab223403 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (ab223403, 1/5000 dilution) for 30 min at 22°C.

Isotype control antibody (black line) was Rabbit IgG (monoclonal) Alexa Fluor<sup>®</sup> 647 (**ab199093**) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.


Acquisition of >5,000 events were collected using a 40 mW Red laser (640nm) and 670/14 bandpass filter.

This antibody gave a positive signal in MCF7 cells fixed with 4% formaldehyde (10 min)/permeabilized with 0.1% PBS-Triton X-100 for 15 min used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-MCM2 antibody [EPR4120] (ab223403)

ab223403 staining MCM2 in MCF7 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab223403 at 1/100 dilution (shown in red) and <a href="mailto:ab195887">ab195887</a>, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). This product also gave a positive signal under the same testing conditions in MCF7 cells fixed with 100% methanol (5 min).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors