abcam

Product datasheet

Alexa Fluor® 647 Anti-STAT3 antibody [EPR787Y] ab300101

Recombinant RabMAb

3 Images

Overview

Product name Alexa Fluor® 647 Anti-STAT3 antibody [EPR787Y]

Description Alexa Fluor® 647 Rabbit monoclonal [EPR787Y] to STAT3

Host species Rabbit

Conjugation Alexa Fluor® 647. Ex: 652nm. Em: 668nm

Tested applications Suitable for: Flow Cyt (Intra), ICC/IF

Species reactivity Reacts with: Human

Immunogen Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

Positive control ICC: HeLa cells. Flow Cyt (Intra): HeLa cells.

General notes

This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or **outlicensing@thermofisher.com**.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, 68% PBS

Purity Protein A purified

Clonality Monoclonal
Clone number EPR787Y

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab300101 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/500.
ICC/IF		1/50.

Target

Function Signal transducer and transcription activator that mediates cellular responses to interleukins,

KITLG/SCF, LEP and other growth factors. Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (PubMed:17344214). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the interleukin-6 (IL-6)-responsive elements identified in the promoters of various acute-phase protein genes. Activated by IL31 through IL31RA. Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed:17344214). Mediates the

effects of LEP on melanocortin production, body energy homeostasis and lactation (By similarity). May play an apoptotic role by transctivating BIRC5 expression under LEP activation

(PubMed:18242580). Cytoplasmic STAT3 represses macroautophagy by inhibiting

EIF2AK2/PKR activity.

Tissue specificity Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.

Involvement in disease Hyperimmunoglobulin E recurrent infection syndrome, autosomal dominant

Autoimmune disease, multisystem, infantile-onset

Sequence similaritiesBelongs to the transcription factor STAT family.

Contains 1 SH2 domain.


Post-translational modifications

Tyrosine phosphorylated upon stimulation with EGF. Tyrosine phosphorylated in response to constitutively activated FGFR1, FGFR2, FGFR3 and FGFR4 (By similarity). Activated through tyrosine phosphorylation by BMX. Tyrosine phosphorylated in response to IL6, IL11, LIF, CNTF, KITLG/SCF, CSF1, EGF, PDGF, IFN-alpha, LEP and OSM. Activated KIT promotes phosphorylation on tyrosine residues and subsequent translocation to the nucleus. Phosphorylated on serine upon DNA damage, probably by ATM or ATR. Serine phosphorylation is important for the formation of stable DNA-binding STAT3 homodimers and maximal transcriptional activity. ARL2BP may participate in keeping the phosphorylated state of STAT3 within the nucleus. Upon LPS challenge, phosphorylated within the nucleus by IRAK1. Upon erythropoietin treatment, phosphorylated on Ser-727 by RPS6KA5. Phosphorylation at Tyr-705 by PTK6 or FER leads to an increase of its transcriptional activity. Dephosphorylation on tyrosine residues by PTPN2 negatively regulates IL6/interleukin-6 signaling.

Cellular localization

Cytoplasm. Nucleus. Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4. Constitutive nuclear presence is independent of tyrosine phosphorylation. Predominantly present in the cytoplasm without stimuli. Upon leukemia inhibitory factor (LIF) stimulation, accumulates in the nucleus. The complex composed of BART and ARL2 plays an important role in the nuclear translocation and retention of STAT3. Identified in a complex with LYN and PAG1.


Images




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-STAT3 antibody [EPR787Y] (ab300101)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervix adenocarcinoma epithelial cells) labelling STAT3 with ab300101 at 1/50 dilution (red). Confocal image showing cytoplasmic and nuclear staining in HeLa cells. The nuclear counterstain is DAPI (blue).

Tubulin is detected with <u>ab195887</u> Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor[®] 488) (green) at 1/200 dilution.



Flow Cytometry (Intracellular) - Alexa Fluor® 647 Anti-STAT3 antibody [EPR787Y] (AB300101) Intracellular Flow Cytometry analysis of HeLa (human cervix adenocarcinoma epithelial cells) with ab300101 at a 1/500 dilution labelling STAT3 (Red). Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. Rabbit monoclonal isotype control - Alexa Fluor[®] 647 (Black). Unlabelled control - Cells without incubation with primary or secondary antibodies (Blue).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors