abcam

Product datasheet

Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] ab184705



4 References 7 Images

Overview

Product name Anti-Alpha Skeletal Muscle Actin antibody [EPR18430]

Description Rabbit monoclonal [EPR18430] to Alpha Skeletal Muscle Actin

Host species Rabbit

Tested applications Suitable for: WB, IHC-P

Species reactivity Reacts with: Mouse, Rat, Human

Immunogen Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

Positive control WB: Human, mouse and rat skeletal muscle and heart lysates; Human bladder lysate. IHC-P:

Mouse and rat skeletal muscle tissues.

General notes IHC is recommended for rat and mouse only.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

Purity Protein A purified

Clonality Monoclonal

1

Clone number EPR18430

Isotype IgG

Applications

The Abpromise guarantee

Our Abpromise guarantee covers the use of ab184705 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000. Detects a band of approximately 42 kDa (predicted molecular weight: 42 kDa).
IHC-P		1/100. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target

Function

Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.

Involvement in disease

Defects in ACTA1 are the cause of nemaline myopathy type 3 (NEM3) [MIM:161800]. A form of nemaline myopathy. Nemaline myopathies are muscular disorders characterized by muscle weakness of varying severity and onset, and abnormal thread-or rod-like structures in muscle fibers on histologic examination. The phenotype at histological level is variable. Some patients present areas devoid of oxidative activity containg (cores) within myofibers. Core lesions are unstructured and poorly circumscribed.

Defects in ACTA1 are a cause of myopathy congenital with excess of thin myofilaments (MPCETM) [MIM:161800]. A congenital muscular disorder characterized at histological level by areas of sarcoplasm devoid of normal myofibrils and mitochondria, and replaced with dense masses of thin filaments. Central cores, rods, ragged red fibers, and necrosis are absent. Defects in ACTA1 are a cause of congenital myopathy with fiber-type disproportion (CFTD) [MIM:255310]; also known as congenital fiber-type disproportion myopathy (CFTDM). CFTD is a genetically heterogeneous disorder in which there is relative hypotrophy of type 1 muscle fibers compared to type 2 fibers on skeletal muscle biopsy. However, these findings are not specific and can be found in many different myopathic and neuropathic conditions.


Sequence similarities

Belongs to the actin family.

Cellular localization

Cytoplasm > cytoskeleton.

Images



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Alpha Skeletal Muscle
Actin antibody [EPR18430] (ab184705)


Immunohistochemical analysis of paraffin-embedded Rat skeletal muscle tissue labeling Alpha Skeletal Muscle Actin with ab184705 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Weak cytoplasm staining in rat skeletal muscle is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab97051** at 1/500 dilution.

Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Western blot - Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] (ab184705)

All lanes : Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] (ab184705) at 1/1000 dilution

Lane 1: Mouse brain tissue lysate

Lane 2: Mouse heart tissue lysate

Lane 3: Mouse kidney tissue lysate

Lane 4: Mouse spleen tissue lysate

Lane 5: Rat brain tissue lysate

Lane 6: Rat heart tissue lysate

Lane 7: Rat kidney tissue lysate

Lane 8: Rat spleen tissue lysate

Lane 9: Human brain tissue lysate

Lane 10: Human heart tissue lysate

Lane 11: Human kidney tissue lysate

Lane 12: Human spleen tissue lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Anti-Rabbit $\lg G$ (HRP), specific to the non-reduced form of $\lg G$ at 1/1000 dilution

Predicted band size: 42 kDa

Observed band size: 42 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Alpha Skeletal Muscle
Actin antibody [EPR18430] (ab184705)


Immunohistochemical analysis of paraffin-embedded Mouse skeletal muscle tissue labeling Alpha Skeletal Muscle Actin with ab184705 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Weak cytoplasm staining in mouse skeletal muscle is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab97051** at 1/500 dilution.

Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Western blot - Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] (ab184705) **All lanes :** Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] (ab184705) at 1/1000 dilution

Lane 1 : Human skeletal muscle lysate

Lane 2 : Mouse skeletal muscle lysate


Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Anti-Rabbit lgG (HRP), specific to the non-reduced form of lgG at 1/10000 dilution

Predicted band size: 42 kDa Observed band size: 42 kDa

Exposure time: 5 seconds



Western blot - Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] (ab184705)




Secondary

Anti-Rabbit lgG (HRP), specific to the non-reduced form of lgG at 1/10000 dilution

Predicted band size: 42 kDa **Observed band size:** 42 kDa

Exposure time: 1 second

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] (ab184705)

Anti-Alpha Skeletal Muscle Actin antibody [EPR18430] (ab184705) at 1/1000 dilution + Human bladder lysate at $10 \mu g$


Secondary

Anti-Rabbit lgG (HRP), specific to the non-reduced form of lgG at 1/10000 dilution

Predicted band size: 42 kDa **Observed band size:** 42 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors