# abcam

# Product datasheet

# APC Anti-Transferrin Receptor antibody [MEM-75], prediluted ab64672

Overview

Product name APC Anti-Transferrin Receptor antibody [MEM-75], prediluted

**Description** APC Mouse monoclonal [MEM-75] to Transferrin Receptor, prediluted

Host species Mouse

**Conjugation** APC. Ex: 645nm, Em: 660nm

Specificity ab64672 reacts with CD71 antigen (transferrin receptor), a 95 kDa type II homodimeric

transmembrane glycoprotein expressed on activated B and lymphocytes, macrophages and erythroid precursors; it is lost on resting blood leukocytes. ab64672 does not block binding of

transferrin to the receptor.

Tested applications Suitable for: Flow Cyt

Species reactivity Reacts with: Human

Immunogen Tissue, cells or virus corresponding to Human Transferrin Receptor. NALM-6 human pre-B cell

line

**General notes**The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

**Properties** 

Form Liquid

**Storage instructions** Shipped at 4°C. Store at +4°C.

**Storage buffer** pH: 7.4

Preservative: 0.097% Sodium azide

Constituents: PBS, 0.2% BSA

**Purity** Size exclusion

**Purification notes**The purified antibody (>95% by SDS-PAGE) is conjugated with cross-linked Allophycocyanin

(APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and

adjusted for direct use.

Clonality Monoclonal
Clone number MEM-75
Isotype IgG1

#### **Applications**

The Abpromise guarantee Our Abpromise guarantee covers the use of ab64672 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		

**Application notes** Flow Cyt: Use  $10\mu$ I for  $10^6$  cells or  $100\mu$ I of whole blood.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

## **Target**

Function	Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the heditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. Positively regulates T and B cell proliferation through iron uptake (PubMed:26642240).  (Microbial infection) Acts as a receptor for new-world arenaviruses: Guanarito, Junin and Machupo virus.	
Involvement in disease	Immunodeficiency 46	
Sequence similarities	Belongs to the peptidase M28 family. M28B subfamily.  Contains 1 PA (protease associated) domain.	
Post-translational modifications	N- and O-glycosylated, phosphorylated and palmitoylated. The serum form is only glycosylated. Proteolytically cleaved on Arg-100 to produce the soluble serum form (sTfR). Palmitoylated on both Cys-62 and Cys-67. Cys-62 seems to be the major site of palmitoylation.	
Cellular localization	Secreted and Cell membrane. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors