

Product datasheet

Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free ab240393

Recombinant RabMAb

★★★★★ <u>1 Abreviews</u> 7 Images

Overview		
Product name	Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free	
Description	Rabbit monoclonal [EPR19932] to Aquaporin 3 - BSA and Azide free	
Host species	Rabbit	
Tested applications	Suitable for: IHC-P	
Species reactivity	Reacts with: Human	
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.	
Positive control	IHC-P: Human kidney, skin, bladder cancer, cervix cancer and skin squamous cell cancer tissues. ICC: A431 (human epidermoid carcinoma epithelial cell)	
General notes	ab240393 is the carrier-free version of <u>ab215853</u> .	
	Our <u>carrier-free</u> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.	
	This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.	
	Use our <u>conjugation kits</u> for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.	
	This product is compatible with the Maxpar [®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar [®] is a trademark of Fluidigm Canada Inc.	
	 This product is a recombinant monoclonal antibody, which offers several advantages including: High batch-to-batch consistency and reproducibility Improved sensitivity and specificity Long-term security of supply Animal-free production For more information <u>see here</u>. Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <u>RabMAb[®] patents</u>. 	

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR19932
lsotype	lgG


Applications

The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab240393 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	★ ★ ★ ★ ★ (1)	Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target		
Function	Water channel required to promote glycerol permeability and water transport across cell membranes. Acts as a glycerol transporter in skin and plays an important role in regulating SC (stratum corneum) and epidermal glycerol content. Involved in skin hydration, wound healing, and tumorigenesis. Provides kidney medullary collecting duct with high permeability to water, thereby permitting water to move in the direction of an osmotic gradient. Slightly permeable to urea and may function as a water and urea exit mechanism in antidiuresis in collecting duct cells. It may play an important role in gastrointestinal tract water transport and in glycerol metabolism.	
Tissue specificity	Widely expressed in epithelial cells of kidney (collecting ducts) and airways, in keratinocytes, immature dendritic cells and erythrocytes. Isoform 2 is not detectable in erythrocytes at the protein level.	
Sequence similarities	Belongs to the MIP/aquaporin (TC 1.A.8) family.	
Domain	Aquaporins contain two tandem repeats each containing three membrane-spanning domains and a pore-forming loop with the signature motif Asn-Pro-Ala (NPA).	
Cellular localization	Basolateral cell membrane. In collecting ducts of kidney.	




Immunocytochemistry/ Immunofluorescence - Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free (ab240393) This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab215853</u>).

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized A431 (human epidermoid carcinoma epithelial cell) labelling Aquaporin 3 with <u>ab215853</u> at 1/100 (5.03 µg/ml) dilution (Green). Confocal image showing mainly membranous staining in A431 cell line.

Nuclear DNA was labelled with DAPI (shown in blue). Secondary used was <u>ab150081</u> Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 µg/ml). <u>ab195889</u> Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5µg/ml) dilution (Magenta).

Negative control: U-87 MG

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free (ab240393) Immunohistochemical analysis of paraffin-embedded human kidney tissue labeling Aquaporin 3 with <u>ab215853</u> at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.


Membranous staining on kidney collecting ducts is observed.

Counter stained with Hematoxylin.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab215853</u>).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free (ab240393)



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free (ab240393)

Immunohistochemical analysis of paraffin-embedded human skin tissue labeling Aquaporin 3 with <u>ab215853</u> at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.

Membranous staining on skin squamous cell is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab215853</u>).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemical analysis of paraffin-embedded human bladder cancer tissue labeling Aquaporin 3 with <u>ab215853</u> at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.


Membranous staining on bladder cancer is observed.

Counter stained with Hematoxylin.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab215853**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free (ab240393)



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Aquaporin 3 antibody [EPR19932] - BSA and Azide free (ab240393)

Immunohistochemical analysis of paraffin-embedded human cervix cancer tissue labeling Aquaporin 3 with <u>ab215853</u> at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.

Membranous staining on cervix cancer is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab215853</u>).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemical analysis of paraffin-embedded human skin squamous cell cancer tissue labeling Aquaporin 3 with <u>ab215853</u> at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.


Membranous staining on human skin squamous cell cancer is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab215853**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors