abcam

Product datasheet

Anti-ARHGAP17 antibody ab74454

* ★ ★ ★ ★ ↑ 1 Abreviews 3 References 1 Image

Overview

Product name Anti-ARHGAP17 antibody

Description Rabbit polyclonal to ARHGAP17

Host species Rabbit

Tested applications Suitable for: WB

Species reactivity Reacts with: Human

Immunogen Synthetic peptide corresponding to Human ARHGAP17 aa 300-400 (internal sequence).

Database link: **Q68EM7**

General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 50% Glycerol (glycerin, glycerine), 0.87% Sodium chloride, PBS

Purity Immunogen affinity purified

Clonality Polyclonal

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab74454 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

1

Application	Abreviews	Notes
WB		1/500 - 1/1000. Detects a band of approximately 95 kDa (predicted molecular weight: 95 kDa).

Target


Relevance

ARHGAP17 is a Rho GTPase-activating protein involved in the maintenance of tight junction by regulating the activity of CDC42, thereby playing a central role in apical polarity of epithelial cells. It specifically acts as a GTPase activator for the CDC42 GTPase by converting it to an inactive GDP-bound state. The complex formed with AMOT acts by regulating the uptake of polarity proteins at tight junctions, possibly by deciding whether tight junction transmembrane proteins are recycled back to the plasma membrane or sent elsewhere. RHGAP17 participates in the Ca(2+)-dependent regulation of exocytosis, possibly by catalyzing GTPase activity of Rho family proteins and by inducing the reorganization of the cortical actin filaments. Acts as a GTPase activator in vitro for RAC1.

Cellular localization

Cytoplasm; Cell membrane (tight junction)

Images



All lanes: Anti-ARHGAP17 antibody (ab74454) at 1/500 dilution

Lane 1: extracts from LOVO cells

Lane 2: extracts from LOVO cells with immunizing peptide at 5 µg

Lysates/proteins at 15 µg per lane.

Predicted band size: 95 kDa **Observed band size:** 95 kDa

 $\textbf{Please note:} \ \ \textbf{All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"}$

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

• We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors