


Product datasheet

Anti-ARID1A antibody [EPR13501] - BSA and Azide free ab217154

KO VALIDATED Recombinant RabMAB

[1 References](#) [11 Images](#)

Overview

Product name	Anti-ARID1A antibody [EPR13501] - BSA and Azide free
Description	Rabbit monoclonal [EPR13501] to ARID1A - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, ChIC/CUT&RUN-seq, Flow Cyt (Intra), IHC-P, WB Unsuitable for: ChIP
Species reactivity	Reacts with: Human Predicted to work with: Mouse, Rat 
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
Positive control	IHC-P: Human kidney and human adenocarcinoma of endometrium without ARID1A mutation tissues. ICC/IF: Wildtype HAP1 and SH-SY5Y cells. WB: HEK-293T and SH-SY5Y cell lysates. ChIC/CUT&RUN-Seq: HCT116 cells.
General notes	<p>ab217154 is the carrier-free version of ab182560.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.20 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR13501
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab217154 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		Use at an assay dependent concentration.
ChIC/CUT&RUN-seq		Use at an assay dependent concentration.
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG (Low endotoxin, Azide free), is suitable for use as an isotype control with this antibody.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		1/1000. Predicted molecular weight: 242 kDa.

Application notes Is unsuitable for ChIP.

Target

Function Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Binds DNA non-specifically. Also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling

complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth.

Tissue specificity

Highly expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, and PBL, and at a much lower level in heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas.

Sequence similarities

Contains 1 ARID domain.


Post-translational modifications

Phosphorylated upon DNA damage, probably by ATM or ATR.

Cellular localization

Nucleus.

Images




ChIP/CUT&RUN sequencing - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

ChIP/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5×10^5 HCT116 cells and 5 μ g of **ab182560** [EPR13501]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.

Additional screenshots of mapped reads can be downloaded [here](#).

The University of Geneva owns patents relevant to ChIP (Chromatin Immuno-Cleavage) methods.

This data was developed using the same antibody clone in a different buffer formulation (**ab182560**).



Western blot - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

All lanes : Anti-ARID1A antibody [EPR13501] (**ab182560**) at 1/1000 dilution

Lane 1 : Wild-type HEK-293T cell lysate

Lane 2 : ARID1A knockout HEK-293T cell lysate

Lane 3 : SH-SY5Y cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 242 kDa

Observed band size: 270 kDa

This data was developed using the same antibody clone in a different buffer formulation (**ab182560**).

Lanes 1 - 3: Merged signal (red and green). Green - **ab182560** observed at 270 kDa. Red - loading control **ab8245** (Mouse anti-GAPDH antibody [6C5]) observed at 37 kDa.

ab182560 was shown to react with ARID1A in wild-type HEK-293T cells in Western blot with loss of signal observed in ARID1A knockout cell line **ab266189** (ARID1A knockout cell lysate **ab257250**). Wild-type HEK-293T and ARID1A knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3 % milk in TBS-T (0.1 % Tween®) before incubation with **ab182560** and **ab8245** (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 h at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

Immunohistochemical analysis of paraffin embedded Human adenocarcinoma of endometrium without ARID1A mutation (Left image) labeling ARID1A using **ab182560** at 1/1000 dilution. Right image: Right picture: paraffine embedded human adenocarcinoma of endometrium with ARID1A mutation. A Ready to use HRP Polymer for Rabbit IgG (prediluted) was used as secondary. Counterstain: Hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab182560**).


Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

Intracellular Flow Cytometry analysis of SH-SY5Y (human neuroblastoma) cells labeling ARID1A with purified **ab182560** at 1/230 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab182560**).



Western blot - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

Anti-ARID1A antibody [EPR13501] (**ab182560**) at 1/1000 dilution + 293T (Human embryonic kidney epithelial cell) whole cell lysate at 20 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 242 kDa

Observed band size: 130,270 kDa

Exposure time: 20 seconds


This data was developed using the same antibody clone in a

different buffer formulation ([ab182560](#)).

Blocking buffer and concentration: 5% NFDm/TBS

Observed band: 130-270 kDa


ARID1A has many mutations which typically generate truncated proteins that are highly prone to degradation. (PMID: 21614196, PMID: 29486633, PMID: 34429326).



Immunocytochemistry/ Immunofluorescence - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

[ab182560](#) staining ARID1A in wild-type HAP1 cells (top panel) and ARID1A knockout HAP1 cells (bottom panel). The cells were fixed with 4% formaldehyde (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with [ab182560](#) at 1/500 dilution and [ab195889](#) at 1/250 dilution (shown in pseudo colour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) ([ab150081](#)) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab182560](#)).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

This IHC data was generated using the same anti-ARID1A antibody clone, EPR13501, in a different buffer formulation (cat# [ab182560](#)).

Immunohistochemical analysis of paraffin embedded Human kidney tissue (Left image) labeling ARID1A using [ab182560](#) at 1/1000 dilution. Right image: Right picture: paraffine embedded human clear cell carcinoma of kidney with ARID1A mutation. A Ready to use HRP Polymer for Rabbit IgG (prediluted) was used as secondary. Counterstain: Hematoxylin.

Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

Clone EPR13501 (ab217154) has been successfully conjugated by Abcam. This image was generated using Anti-ARID1A antibody [EPR13501] (Alexa Fluor® 647). Please refer to [ab216304](#) for protocol details.

[ab216304](#) staining ARID1A in wild-type HAP1 cells (top panel) and ARID1A knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with [ab216304](#) at 1/500 dilution (shown in red) and [ab195887](#) at 1/250 dilution (shown in green) overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

Clone EPR13501 (ab217154) has been successfully conjugated by Abcam. This image was generated using Anti-ARID1A antibody [EPR13501] (Alexa Fluor® 488). Please refer to [ab216112](#) for protocol details.

[ab216112](#) staining ARID1A in wild-type HAP1 cells (top panel) and ARID1A knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with [ab216112](#) at 1/500 dilution (shown in green) and [ab195889](#) at 1/250 dilution (shown in pseudo colour red) overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunofluorescent analysis of SH-SY5Y cells labeling ARID1A with **ab182560** at 1/500 and Goat anti rabbit IgG(Alexa Fluor®555) at 1/200. Image at the right stained with DAPI.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab182560**).

Immunocytochemistry/ Immunofluorescence - Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

Why choose a recombinant antibody?



Anti-ARID1A antibody [EPR13501] - BSA and Azide free (ab217154)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors