abcam

Product datasheet

Anti-ATP6V0D1/P39 antibody ab155594

pH: 7.00

Polyclonal

lgG

1 Image

Overview		
Product name	Anti-ATP6V0D1/P39 antibody	
Description	Rabbit polyclonal to ATP6V0D1/P39	
Host species	Rabbit	
Tested applications	Suitable for: WB	
Species reactivity	Reacts with: Human	
	Predicted to work with: Mouse, Rat, Cow, Xenopus laevis, Zebrafish, Xenopus tropicalis 🛛 🔺	
Immunogen	Recombinant fragment corresponding to Human ATP6V0D1/P39 aa 1-350. Database link: <u>P61421</u>	
Positive control	NT2D1, IMR32 and U87-MG cell lysates.	
General notes	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.	
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As	
Properties		
Form	Liquid	
Storage instructions	Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.	

Preservative: 0.01% Thimerosal (merthiolate)

Immunogen affinity purified

Constituents: 78.99% PBS, 1% BSA, 20% Glycerol (glycerin, glycerine)

Storage buffer

Purity

Clonality

Isotype


Our <u>Abpromise guarantee</u> covers the use of ab155594 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/3000. Predicted molecular weight: 40 kDa.

Target		
Function	Subunit of the integral membrane V0 complex of vacuolar ATPase. Vacuolar ATPase is responsible for acidifying a variety of intracellular compartments in eukaryotic cells, thus providing most of the energy required for transport processes in the vacuolar system. May play a role in coupling of proton transport and ATP hydrolysis (By similarity). May play a role in cilium biogenesis through regulation of the transport and the localization of proteins to the cilium.	
Tissue specificity	Ubiquitous.	
Sequence similarities	Belongs to the V-ATPase V0D/AC39 subunit family.	
Cellular localization	Membrane. Localizes to centrosome and the base of the cilium.	

Images



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors