abcam

Product datasheet

Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] - BSA and Azide free ab243945



RabMAb

7 Images

Overview

Product name Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] - BSA and Azide free

Description Rabbit monoclonal [EPR16401] to ATP6V1B1 + ATP6V1B2 - BSA and Azide free

Host species Rabbit

Tested applications Suitable for: WB, ICC/IF, IHC-P

Species reactivity Reacts with: Mouse, Rat, Human

Immunogen Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

Positive control WB: JAR and RAW 264.7 cell lysates; Mouse brain, mouse kidney and rat kidney tissue

lysates.IHC-P: Human, mouse and rat kidney tissues.ICC/IF: HEK293 and JAR cells.

General notes ab243945 is the carrier-free version of <u>ab200839</u>.

Our <u>carrier-free</u> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our <u>conjugation kits</u> for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

1

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Protein A purified

ClonalityMonoclonalClone numberEPR16401

Isotype IgG


Applications

The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab243945 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Detects a band of approximately 57 kDa (predicted molecular weight: 57 kDa).
ICC/IF		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Images



Immunocytochemistry/ Immunofluorescence - Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] -BSA and Azide free (ab243945) Immunofluorescent analysis of 100% methanol-fixed, 0.1% Triton X-100 permeabilized JAR (Human placenta choriocarcinoma cell line) cells labeling ATP6V1B1 + ATP6V1B2 with ab200839 at 1/250 dilution, followed by Goat anti-rabbit lgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 dilution (green). Cytoplasmic staining on JAR cell line was observed. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:

-ve control 1: <u>ab200839</u> at 1/250 dilution followed by <u>ab150120</u> (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution. -ve control 2: <u>ab7291</u> (anti-Tubulin mouse mAb) at 1/1000 dilution

followed by <u>ab150077</u> (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab200839)

ab200839 MERGED

DAPI -ve control 1 -ve control 2


Immunocytochemistry/ Immunofluorescence - Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] -BSA and Azide free (ab243945)

Immunofluorescent analysis of 100% methanol-fixed, 0.1% Triton X-100 permeabilized HEK293 (Human embryonic kidney) cells labeling ATP6V1B1 + ATP6V1B2 with ab200839 at 1/250 dilution, followed by Goat anti-rabbit lgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 dilution (green). Cytoplasmic staining on HEK293 cell line was observed. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:

-ve control 1: <u>ab200839</u> at 1/250 dilution followed by <u>ab150120</u> (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
-ve control 2: <u>ab7291</u> (anti-Tubulin mouse mAb) at 1/1000 dilution followed by <u>ab150077</u> (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab200839)




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] - BSA and Azide free (ab243945)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling ATP6V1B1 + ATP6V1B2 with <u>ab200839</u> at 1/5000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) secondary antibody at 1/500 dilution. Cytoplasmic staining on Human kidney tissue is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab200839)

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] - BSA and Azide free (ab243945)

Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue labeling ATP6V1B1 + ATP6V1B2 with <u>ab200839</u> at 1/5000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) secondary antibody at 1/500 dilution. Cytoplasmic staining on mouse kidney tissue is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab200839)

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] - BSA and Azide free (ab243945)

Immunohistochemical analysis of paraffin-embedded Rat kidney tissue labeling ATP6V1B1 + ATP6V1B2 with <u>ab200839</u> at 1/5000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) secondary antibody at 1/500 dilution. Cytoplasmic staining on rat kidney tissue is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab200839)

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401] - BSA and Azide free (ab243945)

Immunohistochemical analysis of paraffin-embedded Human cardiac muscle tissue labeling ATP6V1B1 + ATP6V1B2 with ab200839 at 1/5000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) secondary antibody at 1/500 dilution. Human cardiac muscle tissue represents a negative control for ATP6V1B1 + ATP6V1B2. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab200839)

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.











Success from the Ethical standards first experiment Confirmed specificity

compliant Animal-free production

Anti-ATP6V1B1 + ATP6V1B2 antibody [EPR16401]

- BSA and Azide free (ab243945)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors