# abcam

## Product datasheet

## Biotin Anti-Pan Trk antibody [EPR17341] ab252166

Recombinant RabMAb

## 2 Images

#### Overview

**Product name** Biotin Anti-Pan Trk antibody [EPR17341]

**Description** Biotin Rabbit monoclonal [EPR17341] to Pan Trk

**Host species** Rabbit Conjugation **Biotin** 

Suitable for: IHC-P **Tested applications** Species reactivity Reacts with: Human

**Immunogen** Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

Positive control IHC-P: Human normal cerebral cortex tissue.

**General notes** This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity - Long-term security of supply - Animal-free production

For more information see here.

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb patents**.

#### **Properties**

**Form** Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

**Purity** Protein A purified

Clonality Monoclonal Clone number EPR17341

Isotype lgG

#### **Applications**

#### The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab252166 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
IHC-P		1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

#### **Target**

#### Relevance

Family of neurotrophic tyrosine kinase (NTRK1/2/3) genes which encode TrkA, TrkB and TrkC protein kinases. The three family members are activated by different neurotrophins: TrkA is activated by Nerve growth factor (NGF), TrkB by Brain-derived neurotrophic factor (BDNF) or neurotrophin-4 (NT-4) and TrkC by NT-3. Neurotrophin signalling activates cellular pathways involved in the development and the maturation of the central and peripheral nervous systems through regulation of proliferation, differentiation and survival of sympathetic and nervous neurons. Localization TrkA: Cell membrane. Early endosome membrane. Late endosome membrane. Internalized to endosomes upon binding of NGF or NT-3 and further transported to the cell body via a retrograde axonal transport. Localized at cell membrane and early endosomes before nerve growth factor (NGF) stimulation. Recruited to late endosomes after NGF stimulation. Colocalized with RAPGEF2 at late endosomes (By similarity). TrkB: Membrane. TrkC: Membrane.

#### **Images**




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Biotin Anti-Pan Trk antibody [EPR17341] (ab252166)

IHC image of TrkA + TrkB + TrkC staining in a section of formalinfixed paraffin-embedded normal human cerebral cortex\*.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins, performed on a Leica BOND<sup>™</sup>. The section was then incubated overnight at +4°C with ab252166 at 1/100 dilution. Detection was via an HRP conjugated ABC system and DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset negative control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors