

Product datasheet

Anti-C3b / iC3b antibody [7C12] ab231078

[4 References](#) [3 Images](#)

Overview

Product name	Anti-C3b / iC3b antibody [7C12]
Description	Mouse monoclonal [7C12] to C3b / iC3b
Host species	Mouse
Tested applications	Suitable for: WB, ELISA
Species reactivity	Reacts with: Human
Immunogen	Full length native protein (purified). This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: human serum and plasma lysates, human liver whole tissue lysate
General notes	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine
Purity	Protein G purified
Clonality	Monoclonal
Clone number	7C12
Isotype	IgG1
Light chain type	kappa

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab231078 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 68 kDa.
ELISA		Use a concentration of 10 µg/ml.

Target

Function

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

Derived from proteolytic degradation of complement C3, C3a anaphylatoxin is a mediator of local inflammatory process. In chronic inflammation, acts as a chemoattractant for neutrophils (By similarity). It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes.

C3-beta-c: Acts as a chemoattractant for neutrophils in chronic inflammation.

Acylation stimulating protein: adipogenic hormone that stimulates triglyceride (TG) synthesis and glucose transport in adipocytes, regulating fat storage and playing a role in postprandial TG clearance. Appears to stimulate TG synthesis via activation of the PLC, MAPK and AKT signaling pathways. Ligand for C5AR2. Promotes the phosphorylation, ARRB2-mediated internalization and recycling of C5AR2 (PubMed:8376604, PubMed:2909530, PubMed:9059512, PubMed:10432298, PubMed:15833747, PubMed:16333141, PubMed:19615750).

Tissue specificity

Plasma. The acylation stimulating protein (ASP) is expressed in adipocytes and released into the plasma during both the fasting and postprandial periods.

Involvement in disease

Complement component 3 deficiency

Macular degeneration, age-related, 9

Hemolytic uremic syndrome atypical 5

Increased levels of C3 and its cleavage product ASP, are associated with obesity, diabetes and coronary heart disease. Short-term endurance training reduces baseline ASP levels and subsequently fat storage.

Sequence similarities

Contains 1 anaphylatoxin-like domain.


Contains 1 NTR domain.

Post-translational modifications

C3b is rapidly split in two positions by factor I and a cofactor to form iC3b (inactivated C3b) and C3f which is released. Then iC3b is slowly cleaved (possibly by factor I) to form C3c (beta chain + alpha' chain fragment 1 + alpha' chain fragment 2), C3dg and C3f. Other proteases produce other fragments such as C3d or C3g. C3a is further processed by carboxypeptidases to release the C-terminal arginine residue generating the acylation stimulating protein (ASP). Levels of ASP are increased in adipocytes in the postprandial period and by insulin and dietary chylomicrons. Phosphorylated by FAM20C in the extracellular medium.


Cellular localization

Secreted.



ELISA - Anti-C3b / iC3b antibody [7C12] (ab231078)

96-well microtitre plates were coated overnight at 4°C with recombinant human C3, C3b, iC3b, and Factor H proteins, in duplicate at a concentration of 1µg/mL. Plates were blocked with 1% BSA in PBS-T (0.1% Tween®) for 1 hour before incubation with a 10-step 4x serial dilution of ab231078 from 10µg/mL for 1 hour at room temperature. Antibody binding was detected with Goat Anti-Mouse IgG H&L (HRP) (**ab6789**) secondary antibody at a 1 in 10000 dilution for 1 hour at room temperature. Plates were incubated with TMB ELISA substrate for 7 minutes prior to being stopped with Stop solution and absorbance measured at 450nm.



Western blot - Anti-C3b / iC3b antibody [7C12] (ab231078)

All lanes : Anti-C3b / iC3b antibody [7C12] (ab231078) at 1 µg/ml

Lane 1 : Human Sera diluted 1/1000 at 10 µl

Lane 2 : Purified C3 Protein at 0.1 µg

Lane 3 : Purified C3b Protein at 0.1 µg


Lane 4 : Purified iC3b Protein at 0.1 µg

Performed under reducing conditions.

Predicted band size: 68 kDa

Observed band size: 110,125,43,78 kDa

This blot was produced using a 4-12% Bis-tris under the MOPS buffer system. The gel was run at 200V for 55 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% milk before being incubated with ab231078 overnight at 4°C at 1ug/ml. Antibody binding was detected using Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed (**ab216772**) at 1/20000 dilution for 1 hour at room temperature before imaging.



All lanes :

Lane 1 : Human serum diluted 1/100 at 5 µl

Lane 2 : Human plasma diluted 1/100 at 5 µl

Lane 3 : Human liver whole tissue lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 68 kDa

Additional bands at: 42 kDa (possible cleavage fragment)

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 55 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% milk before ab231078 and **ab181602** (Rabbit anti GAPDH) were incubated overnight at 4°C at a 1 µg/ml concentration and 1/10000 dilution respectively. Antibody binding was detected using Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors