



Anti-Caspase-3 antibody ab90437

★★★★★ [9 Abreviews](#) [103 References](#) [3 Images](#)

Overview

Product name	Anti-Caspase-3 antibody
Description	Rabbit polyclonal to Caspase-3
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P
Species reactivity	Reacts with: Human, Saccharomyces cerevisiae
Immunogen	<p>Recombinant full length protein corresponding to Human Caspase-3 aa 1-277.</p> <p>Sequence:</p> <p>MENTENSVDSKSIKNLEPKIIHGSESMDSGISLDNSYKMDY PEMGLCIII NNKNFHKSTGMTSRSGTDVDAANLRETFRNLKYEVRNKN DLTREEVELM RDVSKEDHSKRSSFVCVLLSHGEEGIIFGTNGPVDLKKITN FFRGDRCRS LTGKPKLFIQACRGTELDGCIETDSGVDDDMACHKIPVEA DFLYAYSTA PGYYSWRNSKDGSWFIQSLCAMLKQYADKLEFMHILTRVN RKVATEFESF SFDATFHAKKQIPCIVSMLTKELYFYH</p> <p>Database link: P42574</p> <p> Run BLAST with  Run BLAST with</p>
Positive control	Jurkat whole cell lysate (ab7899), Jurkat cells treated with staurosporine (lysate), HeLa whole cell lysate (ab150035)
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.09% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine)
Purity	Protein A purified
Purification notes	Affinity Purification
Clonality	Polyclonal
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab90437 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★☆ (7)	1/1000. Predicted molecular weight: 32 kDa. A lower molecular weight band refers to a proteolytic cleavage product of caspase-3.
IHC-P		Use at an assay dependent concentration.

Target

Function	Involved in the activation cascade of caspases responsible for apoptosis execution. At the onset of apoptosis it proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-Gly-217' bond. Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Cleaves and activates caspase-6, -7 and -9. Involved in the cleavage of huntingtin.
Tissue specificity	Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.
Sequence similarities	Belongs to the peptidase C14A family.
Post-translational modifications	Cleavage by granzyme B, caspase-6, caspase-8 and caspase-10 generates the two active subunits. Additional processing of the propeptides is likely due to the autocatalytic activity of the activated protease. Active heterodimers between the small subunit of caspase-7 protease and the large subunit of caspase-3 also occur and vice versa. S-nitrosylated on its catalytic site cysteine in unstimulated human cell lines and denitrosylated upon activation of the Fas apoptotic pathway, associated with an increase in intracellular caspase activity. Fas therefore activates caspase-3 not only by inducing the cleavage of the caspase zymogen to its active subunits, but also by stimulating the denitrosylation of its active site thiol.
Cellular localization	Cytoplasm.

Images



Western blot - Anti-Caspase-3 antibody (ab90437)

This image is courtesy of an anonymous abreview.

All lanes : Anti-Caspase-3 antibody (ab90437) at 1/1000 dilution

Lane 1 : Mouse hepatocytes whole cell lysate poly(I:C) treated - 0 hours

Lane 2 : Mouse hepatocytes whole cell lysate poly(I:C) treated - 12 hours


Developed using the ECL technique.

Predicted band size: 32 kDa

Observed band size: 32 kDa


Exposure time: 20 seconds

Blocking with 5% milk for 1 hour at 23°C.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Caspase-3 antibody (ab90437)

Immunohistochemistry analysis of human spleen tissue stained with Caspase-3, pAb at 10µg/ml.



Western blot - Anti-Caspase-3 antibody (ab90437)

All lanes : Anti-Caspase-3 antibody (ab90437) at 1/1000 dilution

Lane 1 : Jurkat cell lysate

Lane 2 : Jurkat cells treated with staurosporine

Lane 3 : MCF-7 cell lysate (negative control)

Developed using the ECL technique.

Predicted band size: 32 kDa

Additional bands at: ~18 kDa (possible cleavage fragment), ~20 kDa (possible cleavage fragment)

Western blot analysis of Caspase-3 pAb.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors