

Product datasheet

Anti-Catalase antibody [EP1929Y] - Peroxisome Marker ab76024

KO VALIDATED Recombinant RabMAb

★★★★★ [1 Abreviews](#) [18 References](#) [7 Images](#)

Overview

Product name	Anti-Catalase antibody [EP1929Y] - Peroxisome Marker
Description	Rabbit monoclonal [EP1929Y] to Catalase - Peroxisome Marker
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, WB, IHC-P Unsuitable for: Flow Cyt or IP
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide within Human Catalase aa 400-500 (C terminal). The exact sequence is proprietary. Database link: P04040 (Peptide available as ab225865)
Positive control	WB: HeLa cell lysate. IHC-P: human brain tissue, human bladder cancer tissue. ICC/IF: HeLa cells.
General notes	Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents . Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	pH: 7.20 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
Purity	Tissue culture supernatant
Clonality	Monoclonal
Clone number	EP1929Y

Isotype

IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab76024 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/100 - 1/250.
WB	★★★★★ (1)	1/10000 - 1/20000. Detects a band of approximately 60 kDa (predicted molecular weight: 60 kDa).
IHC-P		1/100 - 1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Application notes

Is unsuitable for Flow Cyt or IP.

Target

Function

Occurs in almost all aerobically respiring organisms and serves to protect cells from the toxic effects of hydrogen peroxide. Promotes growth of cells including T-cells, B-cells, myeloid leukemia cells, melanoma cells, mastocytoma cells and normal and transformed fibroblast cells.

Involvement in disease

Defects in CAT are the cause of acatalasia (ACATLAS) [MIM:115500]; also known as acatalasemia. This disease is characterized by absence of catalase activity in red cells and is often associated with ulcerating oral lesions.

Sequence similarities

Belongs to the catalase family.


Post-translational modifications

The N-terminus is blocked.

Cellular localization

Peroxisome.

Images



Western blot - Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)

All lanes : Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024) at 1/10000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : CAT knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 60 kDa

Observed band size: 60 kDa

Lanes 1-2: Merged signal (red and green). Green - ab76024 observed at 60 kDa. Red - loading control **ab8245** observed at 37 kDa.


ab76024 Anti-Catalase antibody [EP1929Y] - Peroxisome Marker was shown to specifically react with Catalase in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265250** (knockout cell lysate **ab256859**) was used. Wild-type and Catalase knockout samples were subjected to SDS-PAGE. ab76024 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 10000 and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)

Immunohistochemical analysis of Paraffin-embedded human bladder cancer tissue sections labeling Catalase with ab76024 at 1/1000. Goat Anti-Rabbit IgG H&L (HRP) was used as the secondary antibody. Sections were counterstained with Hematoxylin. Antigen retrieval was heat mediated using [ab93684](#) (Tris/EDTA buffer, pH 9.0).

Granular cytoplasmic staining on human bladder cancer.



Western blot - Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)

All lanes : Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)

Lane 1 : Wild-type HAP1 whole cell lysate

Lane 2 : CAT knockout HAP1 whole cell lysate


Lane 3 : HeLa whole cell lysate

Lysates/proteins at 20 µg per lane.

Predicted band size: 60 kDa

Lanes 1 -3: Merged signal (red and green). Green - ab76024 observed at 60 kDa. Red - loading control, [ab9484](#), observed at 37 kDa.

ab76024 was shown to specifically react with CAT when CAT knockout samples were used. Wild-type and CAT knockout samples were subjected to SDS-PAGE. Ab76024 and [ab9484](#) (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1/5000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)

Immunocytochemistry/Immunofluorescence analysis of HeLa (human cervix adenocarcinoma epithelial cell) labeling Catalase ab76024 at 1/100. Cells were fixed with 100% Methanol.

ab150077, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

ab195889, Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/100 was used as counterstain antibody.

Confocal image showing membranous staining in HeLa cells.



Western blot - Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)


Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024) at 1/20000 dilution + HeLa cell lysate at 10 µg

Secondary

HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 60 kDa

Observed band size: 60 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)

Immunohistochemical staining of Catalase in paraffin embedded human normal brain tissue using ab76024 at a 1/100 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Catalase antibody [EP1929Y] - Peroxisome Marker (ab76024)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors