

Product datasheet

Anti-CCL27 antibody [EPR8582(2)] - BSA and Azide free ab250878

Recombinant RabMAb

4 Images

| Overview | |
|---------------------|---|
| Product name | Anti-CCL27 antibody [EPR8582(2)] - BSA and Azide free |
| Description | Rabbit monoclonal [EPR8582(2)] to CCL27 - BSA and Azide free |
| Host species | Rabbit |
| Tested applications | Suitable for: Flow Cyt (Intra), WB, ICC/IF |
| Species reactivity | Reacts with: Human |
| | Predicted to work with: Mouse, Rat 🛛 🔺 |
| Immunogen | Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. |
| General notes | ab250878 is the carrier-free version of <u>ab186421</u> . |
| | Our <u>carrier-free</u> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency. |
| | This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications. |
| | Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold. |
| | This product is compatible with the Maxpar [®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. $Maxpar^{\mathbb{R}}$ is a trademark of Fluidigm Canada Inc. |
| | This product is a recombinant monoclonal antibody, which offers several advantages including: High batch-to-batch consistency and reproducibility Improved sensitivity and specificity Long-term security of supply Animal-free production For more information <u>see here</u>. Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <u>RabMAb[®] patents</u>. |
| | |

Properties

| Form | Liquid |
|----------------------|---|
| Storage instructions | Shipped at 4°C. Store at +4°C. Do Not Freeze. |
| Storage buffer | pH: 7.2 Constituent: PBS |
| Carrier free | Yes |
| Purity | Protein A purified |
| Clonality | Monoclonal |
| Clone number | EPR8582(2) |
| lsotype | lgG |

Applications


The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab250878 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|------------------|-----------|---|
| Flow Cyt (Intra) | | Use at an assay dependent concentration. |
| WB | | Use at an assay dependent concentration. Detects a band of approximately 10 kDa (predicted molecular weight: 13 kDa). |
| ICC/IF | | Use at an assay dependent concentration. |


| Target | |
|-----------------------|---|
| Function | Chemotactic factor that attracts skin-associated memory T-lymphocytes. May play a role in mediating homing of lymphocytes to cutaneous sites. Binds to CCR10. |
| Tissue specificity | Testis, thymus, placenta, ovary and skin. |
| Sequence similarities | Belongs to the intercrine beta (chemokine CC) family. |
| Cellular localization | Secreted. |
| | |

Images



This data was developed using <u>ab186421</u>, the same antibody clone in a different buffer formulation.Immunofluorescent analysis of 4% paraformaldehyde-fixed A431 cells labeling CCL27 with <u>ab186421</u> at 1/250 dilution followed by Goat anti rabbit lgG (Alexa Fluor® 555) secondary antibody at 1/200 dilution. Counter stained with Dapi.

Immunocytochemistry/ Immunofluorescence - Anti-CCL27 antibody [EPR8582(2)] - BSA and Azide free (ab250878)




Western blot - Anti-CCL27 antibody [EPR8582(2)] -BSA and Azide free (ab250878) Anti-CCL27 antibody [EPR8582(2)] (**ab186421**) at 1/5000 dilution + Human CCL27 recombinant protein at 0.01 µg

Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 13 kDa

This data was developed using <u>ab186421</u>, the same antibody clone in a different buffer formulation.



Flow Cytometry (Intracellular) - Anti-CCL27 antibody [EPR8582(2)] - BSA and Azide free (ab250878)

Why choose a recombinant antibody? Research with Long-term and confidence scalable supply Consistent and Recombinant reproducible results technology Success from the Ethical standards first experiment compliant Confirmed Animal-free specificity production Anti-CCL27 antibody [EPR8582(2)] - BSA and Azide

free (ab250878)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

This data was developed using <u>ab186421</u>, the same antibody clone in a different buffer formulation.

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed HACAT cells labeling CCL27 with <u>ab186421</u> at 1/320 dilution (red) compared to a Rabbit monoclonal IgG isotype control (green), followed by Goat anti rabbit IgG (FITC) secondary antibody at 1/150 dilution. • We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors