abcam

Product datasheet

Anti-CD10 antibody [SP67] - BSA and Azide free ab236235





* ★ ★ ★ ★ ↑ 1 Abreviews 5 Images

Overview

Product name Anti-CD10 antibody [SP67] - BSA and Azide free

Description Rabbit monoclonal [SP67] to CD10 - BSA and Azide free

Host species Rabbit

Tested applications Suitable for: IHC-P, WB, Flow Cyt

Species reactivity Reacts with: Human

Predicted to work with: Mouse, Rat, Chicken, Cow

Synthetic peptide within Human CD10 aa 450-550. The exact sequence is proprietary. **Immunogen**

Database link: P08473

Positive control IHC-P: Human renal cell carcinoma tissue. Flow Cyt: Raji cells.

General notes ab236235 is the carrier-free version of ab227640.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

This product is FOR RESEARCH USE ONLY. For commercial use, please contact

partnerships@abcam.com.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.20

Constituent: PBS

Carrier free Yes

Purity Protein A/G purified

Purification notes Purified from TCS by protein A/G.

Clonality Monoclonal

Clone number SP67
Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab236235 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with 1mM EDTA buffer pH 8.0 before commencing with IHC staining protocol.Primary antibody incubation for 30 minutes at room temperature.
WB		Use at an assay dependent concentration. Detects a band of approximately 86 kDa. Primary antibody incubation for 1 hour at room temperature.
Flow Cyt		Use at an assay dependent concentration.


Function Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids. Biologically important in the destruction of opioid peptides such as Met- and Leuenkephalins by cleavage of a Gly-Phe bond. Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9. Involved in the degradation of atrial natriuretic factor (ANF). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers.

Sequence similarities Belongs to the peptidase M13 family.

Cellular localization Cell membrane.

Images


Target



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD10 antibody [SP67] - BSA and Azide free (ab236235)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human renal cell carcinoma tissue sections labeling CD10 with <u>ab227640</u> at 1/100 dilution (1.89 µg/ml). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 10mins. Rabbit specific IHC polymer detection kit HRP/DAB (<u>ab209101</u>) was used as the secondary antibody. Hematoxylin was used as a counterstain. Positive staining on human renal cell carcinoma, performed on a Leica Biosystems BOND™ RX instrument.


The section was incubated with <u>ab227640</u> for 30 mins at room temperature. This image was generated using <u>ab227640</u>, the same clone, but with a different buffer formulation.



Flow Cytometry - Anti-CD10 antibody [SP67] - BSA and Azide free (ab236235)

Flow Cytometry analysis of Jurkat(Human T cell leukemia T lymphocyte)_negative control cells labeling CD10 with purified ab227640 at 1:200 dilution (1.025 µg/ml) Red. A Goat anti rabbit lgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1:2000 dilution. Isotype control - Rabbit monoclonal lgG (ab172730) / Black. Unlabeled control - Unlabelled cells / Blue.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227642).



Flow Cytometry - Anti-CD10 antibody [SP67] - BSA and Azide free (ab236235)

Flow Cytometry analysis of Raji (Human Burkitt's lymphoma B lymphocyte) cells labeling CD10 with purified ab227640 at 1:200 dilution (1.025 µg/ml) Red. A Goat anti rabbit lgG (Alexa Fluor® 488, ab150081) secondary antibody was used at 1:2000 dilution. Isotype control - Rabbit monoclonal IgG (ab172730) / Black. Unlabeled control - Unlabelled cells / Blue.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227640).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD10 antibody [SP67] -BSA and Azide free (ab236235)

Formalin-fixed, paraffin-embedded human renal cell carcinoma tissue stained for CD10 using ab227640 at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab227640).





Research with confidence Consistent and reproducible results



Long-term and scalable supply Recombinant



Confirmed

specificity





compliant Animal-free production

Anti-CD10 antibody [SP67] - BSA and Azide free (ab236235)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors