abcam

Product datasheet

Anti-CD11b antibody [M1/70] ab8878

Recombinant

★★★★ <u>17 Abreviews</u> <u>138 References</u>

ces 4 Images

Overview

Product name Anti-CD11b antibody [M1/70]

Description Rat monoclonal [M1/70] to CD11b

Host species Rat

Tested applications Suitable for: ICC/IF, Flow Cyt

Unsuitable for: IHC-P

Species reactivity Reacts with: Mouse, Human

ImmunogenTissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.

Positive control Flow Cytometry: RAW 264.7 cells, Human peripheral blood mononuclear cell (PBMC), Mouse

peripheral blood mononuclear cell (PBMC) IF/ICC: RAW 264.7 cells

General notes

This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or

conjugation for your experiments, please contact orders@abcam.com.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

Storage buffer pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)

Purity Protein A purified

Clonality Monoclonal

Clone number M1/70

1

Myeloma NS1 lsotype lgG2b

Light chain type unknown

Applications

The Abpromise guarantee Ou

Our **Abpromise guarantee** covers the use of ab8878 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF	★★★★★ (4)	1/50.
Flow Cyt	★★★★★ (2)	1/1000 - 1/10000. ab18536 - Rat monoclonal lgG2b, is suitable for use as an isotype control with this antibody.

Application notes Is unsuitable for IHC-P.

Target

Function Integrin alpha-M/beta-2 is implicated in various adhesive interactions of monocytes,

macrophages and granulocytes as well as in mediating the uptake of complement-coated particles. It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin alpha-M/beta-2 is also a receptor for fibrinogen, factor X and ICAM1. It recognizes P1 and P2 peptides of fibrinogen

gamma chain.

Tissue specificity Predominantly expressed in monocytes and granulocytes.

Involvement in disease Genetic variations in ITGAM has been associated with susceptibility to systemic lupus

erythematosus type 6 (SLEB6) [MIM:609939]. Systemic lupus erythematosus (SLE) is a chronic, inflammatory and often febrile multisystemic disorder of connective tissue. It affects principally the skin, joints, kidneys and serosal membranes. It is thought to represent a failure of the regulatory

mechanisms of the autoimmune system.

Sequence similarities Belongs to the integrin alpha chain family.


Contains 7 FG-GAP repeats. Contains 1 VWFA domain.

Domain The integrin I-domain (insert) is a VWFA domain. Integrins with I-domains do not undergo


protease cleavage.

Cellular localization Membrane.


Images



Immunocytochemistry/ Immunofluorescence - Anti-CD11b antibody [M1/70] (ab8878)



Negative control: WEHI-231 (PMID: 2457584)




Flow Cytometry - Anti-CD11b antibody [M1/70] (ab8878)

Flow Cytometry analysis of WEHI-231 (Mouse B cell lymphoma B lymphocyte, Left) / RAW 264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage, Right) cells stained for CD11b using AB8878 at a 1/1000 dilution (0.1 μ g) (Red) compared to Rat monoclonal lgG - Isotype Control (Black) and cells without incubation with primary antibody and secondary antibody (Blue). Goat F(ab)2 Anti-Rat lgG Fc (Alexa Fluor® 488, <u>ab150161</u>) at 1/2000 dilution was used as the secondary antibody.


Negative control: WEHI-231.(PMID: 2457584)

Gated on viable cells.



Flow Cytometry - Anti-CD11b antibody [M1/70] (ab8878)

Flow Cytometry analysis of Human peripheral blood mononuclear cell (PBMC) cells labelling CD11b with AB8878 at a 1/10,000 dilution (0.01 µg) (right) compared to Rat monoclonal lgG - lsotype Control (right). Goat F(ab)2 Anti-Rat lgG Fc (Alexa Fluor® 488, ab150161) at 1/2000 dilution was used as the secondary antibody. Gated on viable cells.



Flow Cytometry - Anti-CD11b antibody [M1/70] (ab8878)

Flow Cytometry analysis of Mouse peripheral blood mononuclear cell (PBMC) cells labelling CD11b with AB8878 at a 1/10,000 dilution (0.01 μ g) (right) compared to Rat monoclonal lgG - lsotype Control (right). Goat F(ab)2 Anti-Rat lgG Fc (Alexa Fluor® 488, ab150161) at 1/2000 dilution was used as the secondary antibody. Gated on viable cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors