

Product datasheet

Anti-CD168 antibody [EPR4054] - Low endotoxin, Azide free ab228482

KO VALIDATED Recombinant RabMAB

6 Images

Overview


Product name	Anti-CD168 antibody [EPR4054] - Low endotoxin, Azide free
Description	Rabbit monoclonal [EPR4054] to CD168 - Low endotoxin, Azide free
Host species	Rabbit
Tested applications	Suitable for: WB, IP, IHC-P Unsuitable for: Flow Cyt or ICC/IF
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	Lysates of: MDA-MB-435, LnCaP, T47-D, RAW264.7, PC-12. Human testis tissue
General notes	<p>ab228482 is the carrier-free version of ab124729.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>Our Low endotoxin, azide-free formats have low endotoxin level (≤ 1 EU/ml, determined by the LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.</p>

Properties

Form Liquid

Storage instructions

Shipped at 4°C. Store at +4°C. Do Not Freeze.

Dissociation constant (K_D)K_D = 1.23 x 10⁻¹⁰ M**Storage buffer**pH: 7.20
Constituent: PBS**Carrier free**

Yes

Purity

Protein A purified

Clonality

Monoclonal

Clone number

EPR4054

Isotype

IgG

Applications**The Abpromise guarantee**Our **Abpromise guarantee** covers the use of ab228482 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 84 kDa. The mouse and rat recommendation is based on the WB results. This antibody may not be suitable for IHC with mouse or rat samples.
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. Antigen retrieval is recommended

Application notes

Is unsuitable for Flow Cyt or ICC/IF.

Target**Function**


Involved in cell motility. When hyaluronan binds to HMMR, the phosphorylation of a number of proteins, including the focal adhesion kinase occurs. May also be involved in cellular transformation and metastasis formation, and in regulating extracellular-regulated kinase (ERK) activity.

Tissue specificity

Expressed in breast cancer cell lines and in normal breast tissue.

Cellular localization

Cell surface. Cytoplasm.



Western blot - Anti-CD168 antibody [EPR4054] - Low endotoxin, Azide free (ab228482)

This WB data was generated using the same anti-CD168 antibody clone, EPR4054, in a different buffer formulation (cat# [ab124729](#)).


Lane 1: Wild type HAP1 whole cell lysate (20 µg)

Lane 2: Empty knockout HAP1 whole cell lysate (20 µg)

Lane 3: CD168 whole cell lysate (20 µg)

Lanes 1 - 3: Merged signal (red and green). Green - [ab124729](#) observed at 90 kDa. Red - loading control, [ab9484](#), observed at 37 kDa.

[ab124729](#) was shown to specifically react with CD168 when CD168 knockout samples were used. Wild-type and Empty knockout samples were subjected to SDS-PAGE. [Ab124729](#) and [ab9484](#) (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunoprecipitation - Anti-CD168 antibody [EPR4054] - Low endotoxin, Azide free (ab228482)

[ab124729](#) (purified) at 1:100 dilution (2ug) immunoprecipitating CD168 in MCF-7 (Human breast adenocarcinoma epithelial cell) whole cell lysate.

Lane 1 (input): MCF-7 (Human breast adenocarcinoma epithelial cell) whole cell lysate 10ug


Lane 2 (+): [ab124729](#) & MCF-7 (Human breast adenocarcinoma epithelial cell) whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of [ab124729](#) in MCF-7 (Human breast adenocarcinoma epithelial cell) whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDM/TBST.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab124729](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD168 antibody [EPR4054] - Low endotoxin, Azide free (ab228482)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human gastric carcinoma tissue sections labeling CD168 with purified **ab124729** at 1:250 dilution (7.7 µg/ml). Heat mediated antigen retrieval was performed using EDTA Buffer, pH9.0. Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab124729**).




OI-RD Scanning - Anti-CD168 antibody [EPR4054] - Low endotoxin, Azide free (ab228482)

Equilibrium disassociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab124729**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD168 antibody [EPR4054] - Low endotoxin, Azide free (ab228482)

This IHC data was generated using the same anti-CD168 antibody clone, EPR4054, in a different buffer formulation (cat# **ab124729**).

Immunohistochemical analysis of paraffin-embedded human testis tissue using unpurified **ab124729** at 1/50 - 1/100 dilution

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-CD168 antibody [EPR4054] - Low endotoxin,
Azide free (ab228482)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors