

Product datasheet

Anti-CD3 zeta (phospho Y83) antibody [EP776(2)Y] ab68236

Recombinant RabMAb

[14 References](#) [7 Images](#)

Overview

Product name	Anti-CD3 zeta (phospho Y83) antibody [EP776(2)Y]
Description	Rabbit monoclonal [EP776(2)Y] to CD3 zeta (phospho Y83)
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt (Intra), ICC/IF, Dot blot, WB, IP Unsuitable for: IHC-P
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide within Human CD3 zeta aa 50-150 (phospho Y83). The exact sequence is proprietary. Database link: P20963
Positive control	WB: Jurkat whole cell lysate (ab7899). IP: Jurkat. ICC: Jurkat cells. Flow Cyt (intra): Jurkat cells.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.05% Sodium azide

	Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% PBS
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EP776(2)Y
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab68236 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
Flow Cyt (Intra)		Use at an assay dependent concentration.
ICC/IF		1/100 - 1/250.
Dot blot		Use at an assay dependent concentration.
WB		1/5000 - 1/10000. Detects a band of approximately 18-22 kDa (predicted molecular weight: 18 kDa).
IP		1/50.

Application notes Is unsuitable for IHC-P.

Target


Function	Probable role in assembly and expression of the TCR complex as well as signal transduction upon antigen triggering.
Involvement in disease	Defects in CD247 are the cause of immunodeficiency due to defect in CD3-zeta (CD3ZID) [MIM:610163]. An immunological deficiency characterized by T-cells impaired immune response to alloantigens, tetanus toxoid and mitogens.
Sequence similarities	Belongs to the CD3Z/FCER1G family. Contains 3 ITAM domains.
Domain	The ITAM domains mediate interaction with SHB.
Post-translational modifications	Phosphorylated on Tyr residues after T-cell receptor triggering.
Cellular localization	Membrane.

Images



Flow Cytometry (Intracellular) - Anti-CD3 zeta (phospho Y83) antibody [EP776(2)Y] (ab68236)

Flow Cytometry analysis of Jurkat (human acute T cell leukemia) treated (Red)/untreated (Green) with 1mM pervanadate for 4 hours with purified ab68236 at 1/250 dilution. The secondary antibody was Goat anti rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution. A Rabbit monoclonal IgG (Black) was used as the isotype control and cells without incubation with primary antibody and secondary antibody (Blue) were used as unlabeled control.



Western blot - Anti-CD3 zeta (phospho Y83) antibody [EP776(2)Y] (ab68236)

All lanes : Anti-CD3 zeta (phospho Y83) antibody [EP776(2)Y] (ab68236) at 1/2000 dilution

Lane 1 : Untreated Jurkat cells whole cell lysates

Lane 2 : Jurkat cells were treated with 50mM Pervanadate for 5 minutes whole cell lysates

Lane 3 : Jurkat cells were treated with 50mM Pervanadate for 5 minutes whole cell lysates. Then the membrane was incubated with Alkaline phosphatase.

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 18 kDa

Observed band size: 18 kDa

Exposure time: 3 minutes

Blocking buffer 5% NFDM/TBST

Diluting buffer 5% NFDM/TBST



Western blot - Anti-CD3 zeta (phospho Y83)
antibody [EP776(2)Y] (ab68236)

All lanes : Anti-CD3 zeta (phospho Y83) antibody [EP776(2)Y]
(ab68236) at 1/10000 dilution

Lane 1 : Jurkat cell lysate, untreated.

Lane 2 : Jurkat cell lysate, treated with pervanadate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP-conjugated goat anti-rabbit IgG at 1/2000 dilution

Predicted band size: 18 kDa

Observed band size: 18-22 kDa




Immunocytochemistry/ Immunofluorescence - Anti-
CD3 zeta (phospho Y83) antibody [EP776(2)Y]
(ab68236)

Immunocytochemistry/Immunofluorescence analysis of Jurkat cells (untreated, Per treated and Per+LP treated) labelling CD3 zeta (phospho Y83) with ab68236 (left) and CD3 zeta with **ab40804** (right) both at a dilution of 1/200. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/1000) were also used.

The image shows increased cytoplasmic staining after Pervanadate (1 mM, 30 min) treatment on Jurkat cells. The LP treatment decreased the cytoplasmic staining caused by Pervanadate.

ab40804 was used as a Pan control for ab68236. The results showed cytoplasmic staining on untreated, pervanadate (1 mM, 30 min) treated and Per+LP treated Jurkat cells.



Immunoprecipitation - Anti-CD3 zeta (phospho Y83)
antibody [EP776(2)Y] (ab68236)


CD3 zeta was immunoprecipitated from 0.35 mg Jurkat (Human T cell leukemia T lymphocyte) treated with pervanadate (50mM 5min) whole cell lysate 10 µg with ab68236 at 1/30 dilution (2µg) .
VeriBlot for IP Detection Reagent (HRP)([ab131366](#)) was used at 1/5000 dilution.

Lane 1: Jurkat (Human T cell leukemia T lymphocyte) treated with pervanadate (50mM 5min) whole cell lysate 10 µg

Lane 2: ab68236 IP in Jurkat treated with pervanadate (50mM 5min) whole cell lysate

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab68236 in Jurkat treated with pervanadate (50mM 5min) whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.




Dot Blot - Anti-CD3 zeta (phospho Y83) antibody
[EP776(2)Y] (ab68236)

Dot blot analysis of CD3 zeta (pY83) phospho peptide (lane 1) and CD3 zeta non-phospho peptide (lane 2) labelling CD3 zeta (phospho Y83) with ab68236 at a dilution of 1/1000. A peroxidase-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/2500).

Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: 3 minutes.

Why choose a recombinant antibody?



Anti-CD3 zeta (phospho Y83) antibody [EP776(2)Y]
(ab68236)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors