

## Product datasheet

# Anti-CD36 antibody [EPR22512-58] ab252923

Recombinant RabMAb

★★★★★ [1 Abreviews](#) [3 References](#) [6 Images](#)

### Overview

<b>Product name</b>	Anti-CD36 antibody [EPR22512-58]
<b>Description</b>	Rabbit monoclonal [EPR22512-58] to CD36
<b>Host species</b>	Rabbit
<b>Specificity</b>	The mouse and rat recommendation are based on the IHC results. We do not guarantee WB for mouse and rat.
<b>Tested applications</b>	<b>Suitable for:</b> WB, IHC-P <b>Unsuitable for:</b> Flow Cyt, ICC/IF or IP
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
<b>Positive control</b>	WB: U937 whole cell lysate; human placenta, lung and breast cancer tissue lysates. IHC-P: Human liver tissue; mouse spleen tissue; rat stomach tissue.
<b>General notes</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"><li>- High batch-to-batch consistency and reproducibility</li><li>- Improved sensitivity and specificity</li><li>- Long-term security of supply</li><li>- Animal-free production</li></ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 0.05% BSA, 40% Glycerol
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal

Clone number                      EPR22512-58  
Isotype                                IgG

## Applications

**The Abpromise guarantee**            Our **Abpromise guarantee** covers the use of ab252923 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000. Detects a band of approximately 70-110 kDa (predicted molecular weight: 53 kDa). The mouse and rat recommendation are based on the IHC results. We do not guarantee WB for mouse and rat.
IHC-P	★★★★★ (1)	1/250. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

**Application notes**                      Is unsuitable for Flow Cyt, ICC/IF or IP.

## Target

**Function**                                      Multifunctional glycoprotein that acts as receptor for a broad range of ligands. Ligands can be of proteinaceous nature like thrombospondin, fibronectin, collagen or amyloid-beta as well as of lipidic nature such as oxidized low-density lipoprotein (oxLDL), anionic phospholipids, long-chain fatty acids and bacterial diacylated lipopeptides. They are generally multivalent and can therefore engage multiple receptors simultaneously, the resulting formation of CD36 clusters initiates signal transduction and internalization of receptor-ligand complexes. The dependency on coreceptor signaling is strongly ligand specific. Cellular responses to these ligands are involved in angiogenesis, inflammatory response, fatty acid metabolism, taste and dietary fat processing in the intestine (Probable). Binds long-chain fatty acids and facilitates their transport into cells, thus participating in muscle lipid utilization, adipose energy storage, and gut fat absorption (By similarity) (PubMed:18353783, PubMed:21610069). In the small intestine, plays a role in proximal absorption of dietary fatty acid and cholesterol for optimal chylomicron formation, possibly through the activation of MAPK1/3 (ERK1/2) signaling pathway (By similarity) (PubMed:18753675). Involved in oral fat perception and preferences (PubMed:22240721, PubMed:25822988). Detection into the tongue of long-chain fatty acids leads to a rapid and sustained rise in flux and protein content of pancreatobiliary secretions (By similarity). In taste receptor cells, mediates the induction of an increase in intracellular calcium levels by long-chain fatty acids, leading to the activation of the gustatory neurons in the nucleus of the solitary tract (By similarity). Important factor in both ventromedial hypothalamus neuronal sensing of long-chain fatty acid and the regulation of energy and glucose homeostasis (By similarity). Receptor for thrombospondins, THBS1 and THBS2, mediating their antiangiogenic effects (By similarity). As a coreceptor for TLR4:TLR6 heterodimer, promotes inflammation in monocytes/macrophages. Upon ligand binding, such as oxLDL or amyloid-beta 42, interacts with the heterodimer TLR4:TLR6, the complex is internalized and triggers inflammatory response, leading to NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway, as well as IL1B secretion, through the priming and activation of the NLRP3 inflammasome (By similarity) (PubMed:20037584). Selective and

nonredundant sensor of microbial diacylated lipopeptide that signal via TLR2:TLR6 heterodimer, this cluster triggers signaling from the cell surface, leading to the NF-kappa-B-dependent production of TNF, via MYD88 signaling pathway and subsequently is targeted to the Golgi in a lipid-raft dependent pathway (By similarity) (PubMed:16880211).

(Microbial infection) Directly mediates cytoadherence of Plasmodium falciparum parasitized erythrocytes and the internalization of particles independently of TLR signaling.

#### Involvement in disease

Platelet glycoprotein IV deficiency  
Coronary heart disease 7

#### Sequence similarities

Belongs to the CD36 family.

#### Post-translational modifications

N-glycosylated and O-glycosylated with a ratio of 2:1.


Ubiquitinated at Lys-469 and Lys-472. Ubiquitination is induced by fatty acids such as oleic acid and leads to degradation by the proteasome (PubMed:21610069, PubMed:18353783).

Ubiquitination and degradation are inhibited by insulin which blocks the effect of fatty acids (PubMed:18353783).

#### Cellular localization

Cell membrane. Membrane raft. Golgi apparatus. Apical cell membrane. Upon ligand-binding, internalized through dynamin-dependent endocytosis.

## Images



**All lanes :** Anti-CD36 antibody [EPR22512-58] (ab252923) at 1/1000 dilution

**Lane 1 :** Human placenta tissue lysate

**Lane 2 :** Human lung tissue lysate

**Lane 3 :** Human breast cancer tissue lysate

Lysates/proteins at 20 µg per lane.

#### Secondary

**All lanes :** VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) at 1/1000 dilution

**Predicted band size:** 53 kDa


**Observed band size:** 70-110 kDa

Blocking and dilution buffer: 5% NFDm/TBST.

Exposure times.

Lane 1: 5.5 seconds; Lane 2: 15 seconds; Lane 3: 48 seconds.

The molecular weight observed is due to glycosylation and consistent with what has been described in the literature (PMID:8798390,15677505,20637247).



Western blot - Anti-CD36 antibody [EPR22512-58] (ab252923)

**All lanes** : Anti-CD36 antibody [EPR22512-58] (ab252923) at 1/1000 dilution

**Lane 1** : U937 (human histiocytic lymphoma cell line) whole cell lysate

**Lane 2** : Ramos (human Burkitt's lymphoma cell line) whole cell lysate

**Lane 3** : Jurkat (human T cell leukemia cell line from peripheral blood) whole cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes** : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 100000 cells

**Predicted band size:** 53 kDa


**Observed band size:** 70-110 kDa

**Exposure time:** 58 seconds

Blocking and dilution buffer: 5% NFDM/TBST.

**Negative control:** Ramos, Jurkat (PMID: 27956597, 10744776).

The molecular weight observed is due to glycosylation and consistent with what has been described in the literature (PMID:8798390,15677505,20637247, 8631821).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD36 antibody [EPR22512-58] (ab252923)

Immunohistochemical analysis of paraffin-embedded rat stomach tissue labeling CD36 with ab252923 at 1/250 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on endothelial cells of rat stomach (PMID:30047927, 22777768) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD36 antibody [EPR22512-58] (ab252923)

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labeling CD36 with ab252923 at 1/250 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on endothelial cells of mouse spleen (PMID:30047927, 22777768) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue labeling CD36 with ab252923 at 1/250 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on endothelial cells of human liver (PMID:30047927, 22777768) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD36 antibody  
[EPR22512-58] (ab252923)

Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-CD36 antibody [EPR22512-58] (ab252923)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors