

## Product datasheet

# Anti-CD45RA antibody [MRC OX-33] - BSA and Azide free ab244569

[1 References](#) [2 Images](#)

### Overview

<b>Product name</b>	Anti-CD45RA antibody [MRC OX-33] - BSA and Azide free
<b>Description</b>	Mouse monoclonal [MRC OX-33] to CD45RA - BSA and Azide free
<b>Host species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt, IHC-Fr
<b>Species reactivity</b>	<b>Reacts with:</b> Rat, Human
<b>Immunogen</b>	Full length native protein (purified) corresponding to Rat CD45RA. Purified from spleen.
<b>Positive control</b>	Flow Cyt: Lewis rat splenocytes. IHC-Fr: Rat Spleen
<b>General notes</b>	<p>ab244569 is the carrier-free version of <a href="#">ab33933</a>.</p> <p>Spleen cells from immunised BALB/c mice were fused with cells of the NSO/U myeloma cell line. This clone has been described reacting with paraffin embedded material following PLP fixation (see Whiteland et al., 1995). It only labels B cells among thoracic duct lymphocytes, with little labelling in bone marrow and none on thymocytes (Barclay et al., 1987).</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p>

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

## Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at +4°C. Do Not Freeze.
<b>Storage buffer</b>	Constituent: PBS
<b>Carrier free</b>	Yes
<b>Purity</b>	Protein G purified
<b>Purification notes</b>	Purified from TCS.
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	MRC OX-33
<b>Isotype</b>	IgG1
<b>Light chain type</b>	kappa

## Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab244569 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use a concentration of 0.2 µg/ml.
IHC-Fr		Use a concentration of 1 µg/ml.

## Target

<b>Function</b>	Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity.
<b>Involvement in disease</b>	Severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive Multiple sclerosis
<b>Sequence similarities</b>	Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily. Contains 2 fibronectin type-III domains. Contains 2 tyrosine-protein phosphatase domains.
<b>Domain</b>	The first PTPase domain interacts with SKAP1.


**Post-translational  
modifications**

Heavily N- and O-glycosylated.

**Cellular localization**

Membrane. Membrane raft. Colocalized with DPP4 in membrane rafts.

**Images**




Lewis rat splenocytes stained with **ab33933** (right) or mouse IgG1k (**ab170190**) isotype (left). Lewis rat splenocytes were incubated for 30 min on ice in 1x PBS / 10 % rat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (**ab33933**) or mouse IgG1k isotype (**ab170190**) ( $1 \times 10^6$  in 100  $\mu$ l at 0.2  $\mu$ g/ml) for 30 min on ice.

The secondary antibody Goat anti-mouse IgG H&L (Alexa Fluor® 488, pre-adsorbed) (**ab150177**) was used at 1/2000 dilution for 30 min at 4°C. The cells were simultaneously stained with CD3 antibody.

Acquisition of >30,000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter. Events were gated on viable lymphocytes.

This data was developed using the same antibody clone in a different buffer formulation containing PBS and sodium azide (**ab33933**).



This data was developed using the same antibody clone in a different buffer formulation containing PBS and sodium azide (**ab33933**).

IHC image of CD45RA staining in a section of frozen normal Rat Spleen.

The section was fixed using 10% formaldehyde in 1XPBS for 10 minutes. No antigen retrieval step was performed prior to staining. Non-specific protein-protein interactions were then blocked in TBS containing 0.025% (v/v) Triton X-100, 0.3M glycine and 1% (w/v) BSA for 1h at room temperature. The section was then incubated overnight at +4°C in TBS containing 0.025% (v/v) Triton X-100 and 1% (w/v) BSA with **ab33933** at 1  $\mu$ g/ml. The section was then incubated with **ab150117** (Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preabsorbed, (Shown in green) 1/1000) for 1 hour at room temperature. DAPI was used to stain the cell nuclei (blue). The secondary-only control insert image is taken from an identical assay without primary antibody. The section was then mounted using Fluoromount®.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

For IHC staining systems (automated and non-automated), customers should optimize variable parameters such as antibody concentrations and incubation times.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors