

Product datasheet

Anti-CD47 antibody [CD47/2937] - BSA and Azide free ab260480

[4 Images](#)

Overview

Product name	Anti-CD47 antibody [CD47/2937] - BSA and Azide free
Description	Mouse monoclonal [CD47/2937] to CD47 - BSA and Azide free
Host species	Mouse
Tested applications	Suitable for: IHC-P, WB, Protein Array
Species reactivity	Reacts with: Human
Immunogen	Recombinant fragment within Human CD47 aa 18-135. The exact sequence is proprietary. Database link: Q08722
Positive control	WB: U-87 and THP-1 cell lysates. IHC-P: Human prostate carcinoma tissue.
General notes	ab260480 is the carrier-free version of ab260418 .

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

Properties

Form	Liquid
-------------	--------

Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A/G purified
Purification notes	Purified from bioreactor concentrate.
Clonality	Monoclonal
Clone number	CD47/2937
Isotype	IgG2c
Light chain type	kappa

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab260480 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use a concentration of 1 - 2 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Primary incubation for 30 minutes at room temperature.
WB		Use a concentration of 1 - 2 µg/ml. Predicted molecular weight: 35 kDa.
Protein Array		Use at an assay dependent concentration.

Target

Function	Has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. Plays an important role in memory formation and synaptic plasticity in the hippocampus (By similarity). Receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation. May play a role in membrane transport and/or integrin dependent signal transduction. May prevent premature elimination of red blood cells. May be involved in membrane permeability changes induced following virus infection.
Tissue specificity	Very broadly distributed on normal adult tissues, as well as ovarian tumors, being especially abundant in some epithelia and the brain.
Sequence similarities	Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Cellular localization	Cell membrane.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD47 antibody [CD47/2937] (ab260480)

Formalin-fixed, paraffin-embedded human prostate carcinoma tissue labeling CD47 using **ab260418** at 2 µg/ml in immunohistochemical analysis.

Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.

This data was developed using the same antibody clone in a different formulation containing BSA and azide (**ab260418**).



Western blot - Anti-CD47 antibody [CD47/2937] (ab260480)


All lanes : Anti-CD47 antibody [CD47/2937] (**ab260418**) at 2 µg/ml

Lane 1 : U-87 (human glioblastoma-astrocytoma epithelial cell line) cell lysate

Lane 2 : THP-1 (human monocytic leukemia cell line) cell lysate

Predicted band size: 35 kDa

This data was developed using the same antibody clone in a different formulation containing BSA and azide (**ab260418**).




Protein Array - Anti-CD47 antibody [CD47/2937]
(ab260480)

Array of >19,000 full-length human proteins, probed with **ab260418**.

Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

This data was developed using the same antibody clone in a different formulation containing BSA and azide (**ab260418**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD47 antibody [CD47/2937] (ab260480)

Formalin-fixed, paraffin-embedded human prostate carcinoma tissue labeling CD47 using **ab260418** at 2 µg/ml in immunohistochemical analysis.

Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.

This data was developed using the same antibody clone in a different formulation containing BSA and azide (**ab260418**).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors