abcam

Product datasheet

Anti-CD63 antibody ab231975

5 References 8 Images

Overview

Product name Anti-CD63 antibody

Description Rabbit polyclonal to CD63

Host species Rabbit

Tested applications

Suitable for: IHC-P, WB

Species reactivity

Reacts with: Human, Pig

Immunogen Recombinant fragment corresponding to Human CD63 aa 100-250. N-terminal His and GST

tags, Expressed in E.coli. Database link: **P08962**

Run BLAST with
Run BLAST with

Positive control WB: Recombinant human CD63 protein; Human serum; Pig liver lysate; A-375 cell lysate. IHC-P:

Human colon cancer, stomach, prostate adenoma, skin cancer, breast cancer and liver tissue.

General notesThe Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer pH: 7.40

Preservative: 0.011% Proclin 300

Constituents: 55.77% Glycerol (glycerin, glycerine), 44.219% PBS

Purity Immunogen affinity purified

Purification notesAntigen-specific affinity chromatography followed by Protein A affinity chromatography.

Clonality Polyclonal

Isotype IgG

1

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab231975 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use a concentration of 5 - 20 µg/ml.
WB		Use a concentration of 0.5 - 2 µg/ml. Predicted molecular weight: 26 kDa.

Target

Function This antigen is associated with early stages of melanoma tumor progression. May play a role in

growth regulation.

Tissue specificity Dysplastic nevi, radial growth phase primary melanomas, hematopoietic cells, tissue


macrophages.

Sequence similaritiesBelongs to the tetraspanin (TM4SF) family.

Cellular localization Cell membrane. Lysosome membrane. Late endosome membrane. Also found in Weibel-Palade

bodies of endothelial cells. Located in platelet dense granules.

Images



All lanes: Anti-CD63 antibody (ab231975) at 2 µg/ml


Lane 1 : Human serum
Lane 2 : Pig liver lysate

Lane 3: A375 (human malignant melanoma cell line) cell lysate

Secondary


All lanes : HRP-Linked Guinea pig anti-Rabbit at 1/2000 dilution

Predicted band size: 26 kDa




Anti-CD63 antibody (ab231975) at 2 $\mu g/ml$ + Recombinant human CD63 protein

Predicted band size: 26 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD63 antibody (ab231975)

Formalin-fixed, paraffin-embedded human liver tissue stained for CD63 using ab231975 at 20 $\mu g/ml$ in immunohistochemical analysis. DAB staining.




Formalin-fixed, paraffin-embedded human breast cancer tissue stained for CD63 using ab231975 at 20 $\mu g/ml$ in immunohistochemical analysis. DAB staining.

Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD63 antibody (ab231975)




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD63 antibody (ab231975)

Formalin-fixed, paraffin-embedded human skin cancer tissue stained for CD63 using ab231975 at 20 μ g/ml in immunohistochemical analysis. DAB staining.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD63 antibody (ab231975)

Formalin-fixed, paraffin-embedded human prostate adenoma tissue stained for CD63 using ab231975 at 20 μ g/ml in immunohistochemical analysis. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD63 antibody (ab231975)

Formalin-fixed, paraffin-embedded human stomach tissue stained for CD63 using ab231975 at 20 μ g/ml in immunohistochemical analysis. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD63 antibody (ab231975)

Formalin-fixed, paraffin-embedded human colorectal cancer tissue stained for CD63 using ab231975 at 20 μ g/ml in immunohistochemical analysis. DAB staining.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors