abcam

Product datasheet

Anti-CXCL7/PBP antibody [EPR20957-7] - BSA and Azide free (Detector) ab245060

Recombinant RabMAb

2 Images

Overview

General notes

Product name Anti-CXCL7/PBP antibody [EPR20957-7] - BSA and Azide free (Detector)

Description Rabbit monoclonal [EPR20957-7] to CXCL7/PBP - BSA and Azide free (Detector)

Host species Rabbit

Tested applications Suitable for: Sandwich ELISA

Species reactivity Reacts with: Mouse

Immunogen Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.

> ab245060 is a BSA and Azide Free antibody supplied in an unconjugated format and it is suitable for sandwich ELISAs to quantify Mouse CXCL7 / PBP. The recommended pair for

sandwich ELISA is:

Capture: ab244787, Mouse CXCL7 / PBP Capture Antibody (unconjugated) Detector: ab245060, Mouse CXCL7 / PBP Detector Antibody (unconjugated)

The reference range value is 2.73 - 175 pg/ml.

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

The recommended antibody orientation is based on internal optimization for ELISA-based assays. Antibody orientation is assay dependent and needs to be optimized for each assay type. Please note that the range provided for this antibody is only an estimation based on the performance of the product using the recommended antibody pair. Performance of the antibody pair will depend on the specific characteristics of your assay. We guarantee the product works in sandwich ELISA, but we do not guarantee the sensitivity or dynamic range of the antibody in your assay.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C.

Storage buffer Constituent: 100% PBS

Carrier free Yes

Purity Protein A purified

Clonality Monoclonal
Clone number EPR20957-7

Isotype IgG

Applications

The Abpromise guarantee

Our Abpromise guarantee covers the use of ab245060 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Sandwich ELISA		Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [EPR20957-97] to CXCL7/PBP - BSA and Azide free (Capture) (ab244787).

Target

Function

LA-PF4 stimulates DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by human synovial cells. NAP-2 is a ligand for CXCR1 and CXCR2, and NAP-2, NAP-2(73), NAP-2(74), NAP-2(1-66), and most potent NAP-2(1-63) are chemoattractants and activators for neutrophils. TC-1 and TC-2 are antibacterial proteins, in vitro released from activated platelet alpha-granules. CTAP-III(1-81) is more potent than CTAP-III desensitize chemokine-induced neutrophil activation.

Sequence similarities

Belongs to the intercrine alpha (chemokine CxC) family.

Post-translational modifications

Proteolytic removal of residues 1-9 produces the active peptide connective tissue-activating peptide III (CTAP-III) (low-affinity platelet factor IV (LA-PF4)).

Proteolytic removal of residues 1-13 produces the active peptide beta-thromboglobulin, which is

released from platelets along with platelet factor 4 and platelet-derived growth factor.

NAP-2(1-66) is produced by proteolytical processing, probably after secretion by leukocytes other


than neutrophils.

NAP-2(73) and NAP-2(74) seem not be produced by proteolytical processing of secreted precursors but are released in an active form from platelets.

Cellular localization


Secreted.

Images



Sandwich ELISA - Anti-CXCL7/PBP antibody
[EPR20957-7] - BSA and Azide free (Detector)
(ab245060)

Representative standard curve from corresponding SimpleStep ELISA® Kit (ab236713).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors