

## Product datasheet

### Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker ab76318

Recombinant RabMAb

★★★★★ [14 Abreviews](#) [106 References](#) [13 Images](#)

#### Overview

<b>Product name</b>	Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker
<b>Description</b>	Rabbit monoclonal [EP1607IHCY] to Cytokeratin 10 - Cytoskeleton Marker
<b>Host species</b>	Rabbit
<b>Specificity</b>	Some customers have successfully used this antibody to staining Cytokeratin 10 in mouse tissue. Please see Abreviews for more details.
<b>Tested applications</b>	<b>Suitable for:</b> ICC/IF, WB, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
<b>Positive control</b>	WB: Hacat and A431 cell lysates, human fetal, rat and mouse skin tissue lysates. IHC-P: Human skin and tonsil tissues and mouse skin tissue. ICC/IF: HACAT cells.
<b>General notes</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol, 0.05% BSA</p>
<b>Purity</b>	Protein A purified

Clonality	Monoclonal
Clone number	EP1607IHCY
Isotype	IgG

## Applications


**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab76318 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF	★★★★★ (1)	1/150. <b>For unpurified use at 1/100 - 1/250.</b>
WB		1/10000. Detects a band of approximately 60 kDa (predicted molecular weight: 60 kDa). <b>For unpurified use at 1/50000 - 1/100000.</b>
IHC-P	★★★★★ (10)	1/2500 - 1/5000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <b>IHC antigen retrieval protocols</b> . <b>For unpurified use at 1/100 - 1/250.</b> <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

## Target


<b>Tissue specificity</b>	Seen in all suprabasal cell layers including stratum corneum.
<b>Involvement in disease</b>	<p>Defects in KRT10 are a cause of bullous congenital ichthyosiform erythroderma (BCIE) [MIM:113800]; also known as epidermolytic hyperkeratosis (EHK) or bullous erythroderma ichthyosiformis congenita of Brocq. BCIE is an autosomal dominant skin disorder characterized by widespread blistering and an ichthyotic erythroderma at birth that persist into adulthood. Histologically there is a diffuse epidermolytic degeneration in the lower spinous layer of the epidermis. Within a few weeks from birth, erythroderma and blister formation diminish and hyperkeratoses develop.</p> <p>Defects in KRT10 are a cause of ichthyosis annular epidermolytic (AEI) [MIM:607602]; also known as cyclic ichthyosis with epidermolytic hyperkeratosis. AEI is a skin disorder resembling bullous congenital ichthyosiform erythroderma. Affected individuals present with bullous ichthyosis in early childhood and hyperkeratotic lichenified plaques in the flexural areas and extensor surfaces at later ages. The feature that distinguishes AEI from BCIE is dramatic episodes of flares of annular polycyclic plaques with scale, which coalesce to involve most of the body surface and can persist for several weeks or even months.</p>
<b>Sequence similarities</b>	Belongs to the intermediate filament family.

## Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 10 antibody [EP1607IHCY] (ab76318)

This image is courtesy of an abreview submitted by Carl Hobbs, King's College London, United Kingdom



Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 10 antibody [EP1607IHCY] (ab76318)



Western blot - Anti-Cytokeratin 10 antibody [EP1607IHCY] (ab76318)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse skin tissue labelling Cytokeratin 10 with unpurified ab76318 at a 1/10000 dilution. The sections were subjected to heat mediated antigen retrieval. The sections were then blocked using 1% BSA for 10 mins at 21°C. ab76318 was incubated for 2 hours at 21°C. A biotin-conjugated goat anti-rabbit IgG polyclonal was used as the secondary antibody (1/250).

Immunocytochemistry/Immunofluorescence analysis of HACAT cells labelling Cytokeratin 10 with purified ab76318 at 1/150. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/500) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.

Control 1: primary antibody (1/100) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).

**All lanes** : Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker (ab76318) at 1/10000 dilution (purified)

**Lane 1** : HACAT cell lysate

**Lane 2** : A431 cell lysate

Lysates/proteins at 20 µg per lane.


**Secondary**

**All lanes** : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

**Predicted band size:** 60 kDa

**Observed band size:** 60 kDa


Blocking and diluting buffer: 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 10 antibody [EP1607IHCY] (ab76318)

This image is courtesy of an abreview submitted by Carl Hobbs, King's College London, United Kingdom

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human skin tissue labelling Cytokeratin 10 with unpurified ab76318 at a 1/6000 dilution. The sections were subjected to heat mediated antigen retrieval. The sections were then blocked using 1% BSA for 10 mins at 21°C. ab76318 was incubated for 2 hours at 21°C. A biotin-conjugated goat anti-rabbit IgG polyclonal was used as the secondary antibody (1/200).



Western blot - Anti-Cytokeratin 10 antibody [EP1607IHCY] (ab76318)

Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker (ab76318) at 1/50000 dilution (purified) + Human fetal skin tissue lysate at 20 µg


### Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

**Predicted band size:** 60 kDa

**Observed band size:** 60 kDa

Blocking and diluting buffer: 5% NFDM/TBST.



Western blot - Anti-Cytokeratin 10 antibody  
[EP1607IHCY] (ab76318)

Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker (ab76318) at 1/50000 dilution (purified) + Rat skin tissue lysate at 20 µg


#### Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

**Predicted band size:** 60 kDa


**Observed band size:** 60 kDa

Blocking and diluting buffer: 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 10 antibody  
[EP1607IHCY] - Cytoskeleton Marker (ab76318)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse skin tissue labelling Cytokeratin 10 with purified ab76318 at 1/5000. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Western blot - Anti-Cytokeratin 10 antibody  
[EP1607IHCY] (ab76318)

Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker (ab76318) at 1/200000 dilution (purified) + Mouse skin tissue lysate at 10  $\mu$ g


#### Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

**Predicted band size:** 60 kDa

**Observed band size:** 60 kDa


Blocking and diluting buffer: 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 10 antibody  
[EP1607IHCY] - Cytoskeleton Marker (ab76318)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human tonsil tissue labelling Cytokeratin 10 with purified ab76318 at 1/5000. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.





Western blot - Anti-Cytokeratin 10 antibody  
[EP1607IHCY] (ab76318)


Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker (ab76318) at 1/100000 dilution + HaCat cell lysate at 10 µg

### Secondary

HRP-conjugated goat anti-rabbit IgG at 1/2000 dilution

**Predicted band size:** 60 kDa

**Observed band size:** 60 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 10 antibody  
[EP1607IHCY] - Cytoskeleton Marker (ab76318)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human normal skin tissue labelling Cytokeratin 10 with unpurified ab76318.

Green - CK10, red - PI.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 10 antibody [EP1607IHCY] (ab76318)

This image is courtesy of an abreview submitted by Carl Hobbs, King's College London, United Kingdom

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat skin tissue labelling Cytokeratin 10 with unpurified ab76318 at a 1/10000 dilution. The sections were subjected to heat mediated antigen retrieval. The sections were then blocked using 1% BSA for 10 mins at 21°C. ab76318 was incubated for 2 hours at 21°C. A biotin-conjugated goat anti-rabbit IgG polyclonal was used as the secondary antibody (1/250).

#### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Cytokeratin 10 antibody [EP1607IHCY] - Cytoskeleton Marker (ab76318)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,



please visit <https://www.abcam.com/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors